

This is a copy of the published version, or version of record, available on the publisher's website. This version does not track changes, errata, or withdrawals on the publisher's site.

Quantum fluctuations in the quasi-one-dimensional non-Fermi liquid system CeCo_2Ga_8 investigated using μSR

A. Bhattacharyya, D. T. Adroja, J. S. Lord, L. Wang, Y. Shi,
K. Panda, H. Luo and A. M. Strydom

Published version information

Citation: A Bhattacharyya et al. 'Quantum fluctuations in the quasi-one-dimensional non-Fermi liquid system CeCo_2Ga_8 investigated using μSR .' Physical Review B, vol. 101, no. 21 (2020): 214437.

DOI: [10.1103/PhysRevB.101.214437](https://doi.org/10.1103/PhysRevB.101.214437)

This version is made available in accordance with publisher policies. Please cite only the published version using the reference above. This is the citation assigned by the publisher at the time of issuing the APV. Please check the publisher's website for any updates.

This item was retrieved from **ePubs**, the Open Access archive of the Science and Technology Facilities Council, UK. Please contact epublications@stfc.ac.uk or go to <http://epubs.stfc.ac.uk/> for further information and policies.

Quantum fluctuations in the quasi-one-dimensional non-Fermi liquid system CeCo_2Ga_8 investigated using μSR

A. Bhattacharyya^{1,*}, D. T. Adroja^{2,3,†}, J. S. Lord², L. Wang^{4,5}, Y. Shi^{4,5,6}, K. Panda,¹
H. Luo,^{4,5,6} and A. M. Strydom³

¹*Department of Physics, Ramakrishna Mission Vivekananda Educational and Research Institute, Belur Math, Howrah 711202, West Bengal, India*

²*ISIS Facility, Rutherford Appleton Laboratory, Chilton, Didcot Oxon OX11 0QX, United Kingdom*

³*Highly Correlated Matter Research Group, Physics Department, University of Johannesburg, P.O. Box 524, Auckland Park 2006, South Africa*

⁴*Beijing National Laboratory for Condensed Matter Physics, Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China*

⁵*School of Physical Sciences, University of Chinese Academy of Sciences, Beijing 100190, China*

⁶*Songshan Lake Materials Laboratory, Dongguan, Guangdong 523808, China*



(Received 20 March 2020; revised manuscript received 10 May 2020; accepted 2 June 2020; published 22 June 2020)

Reduced dimensionality offers a piece of crucial information in deciding the type of the quantum ground state in heavy fermion materials. Here we have examined the stoichiometric CeCo_2Ga_8 compound, which crystallizes in a quasi-one-dimensional crystal structure with Ga-Ce-Co chains along the c axis. The low-temperature behavior of the magnetic susceptibility ($\chi \sim T^{-0.2}$), heat capacity ($C_p/T \sim -\ln T$), and resistivity ($\rho \sim T^n$ with $n \sim 1$) strongly confirms the non-Fermi liquid ground state of CeCo_2Ga_8 . We study the low-energy spin dynamics of the CeCo_2Ga_8 compound utilizing zero-field (ZF) and longitudinal-field muon spin relaxation (μSR) measurements. ZF- μSR measurement reveals the absence of long-range magnetic ordering down to 70 mK. Interestingly, below 1 K the electronic relaxation rate rises sharply, suggesting the appearance of low-energy quantum spin fluctuations in CeCo_2Ga_8 .

DOI: [10.1103/PhysRevB.101.214437](https://doi.org/10.1103/PhysRevB.101.214437)

I. INTRODUCTION

Searching for the quantum critical point (QCP) is a great challenge in strongly correlated materials since it only emerges at zero temperature by varying a control parameter such as the magnetic field, pressure, or chemical doping or alloying [1–3]. Heavy fermion (HF) materials exhibit many exotic states in the vicinity of a magnetic QCP, including non-Fermi liquid (NFL) and unconventional superconductivity [1,4–8]. At the QCP, the quantum fluctuations dominate over the thermal fluctuations that break the predictions of well-known Landau–Fermi liquid behavior [9,10], and hence the system exhibits NFL behavior. The nature of quantum fluctuations and development of magnetic correlations will depend on the dimensionality of the systems, and hence it is very important to investigate the effect of dimensionality on the QCP/NFL. So far, most Ce-based QCP/NFL systems investigated are two-dimensional (2D) or 3D, and there are no reports on 1D Ce-based NFL systems [8,11,12]. Furthermore, the physical properties of Ce-based compounds at low temperatures exhibit Fermi liquid behavior predicted by Landau theory [13]. For example, at low temperatures the electrical resistivity $\rho \sim T^2$, the heat capacity $C \sim T$, and the dc magnetic susceptibility is independent of the temperature

[14–18]. Interestingly some of the Ce- and Yb-based materials deviate from conventional Fermi liquid behavior to so-called NFL behavior, which can be tuned from an antiferromagnetic ground state to a zero-temperature QCP, where quantum fluctuations are responsible for the NFL behavior. In the case of NFL compounds $\rho \sim T^n$ ($1 \leq n < 2$), $C/T \sim -\ln T$ or $C/T \sim a - bT^{1/2}$, and $\chi \sim T^{-p}$ ($p < 1$) [14–18].

To accommodate deeper insight into the specific nature of the QCP, both theoretical and experimental efforts have been made frequently in recent years. Still, most of them focus on the quasi-2D or 3D HF [8,11,12]. Considering that dimensionality is a fundamental component in defining the unique NFL attributes in these materials, a lower dimension predicts a substantial magnetic frustration parameter, and it is imperative to search the QCPs in quasi-1D HF compounds, whose science can be easily approximated by the density matrix renormalization-group method as well as the mean-field approximation [19]. The QCP has been observed in $\text{CeCu}_{6-x}\text{T}_x$ ($T = \text{Au, Ag}$) [20,21], in which the NFL at $x = 0.1$ and an ambient pressure is driven to a magnetically ordered state via further doping in antiferromagnetic ordered HF compounds such as CeIn_3 [22,23] and CePd_2Si_2 [24]. To date, there exist only a few undoped or stoichiometric materials which exhibit NFL states at ambient pressure, such as UBe_{13} [25,26], CeNi_2Ge_2 [27], CeCu_2Si_2 [27], and CeRhBi [28,29].

As chemically disordered NFL states remain challenging to understand by theoretical models, it is highly desirable to

*amitava.bhattacharyya@rkmvu.ac.in

†devashibhai.adroja@stfc.ac.uk

examine stoichiometric and homogeneous systems, which are prototypical materials for theoretical modeling. The recently discovered quasi-1D Kondo lattice system CeCo_2Ga_8 , which crystallizes in the YbCo_2Ga_8 -type orthorhombic structure, provides us a rare opportunity to scrutinize a QCP at ambient pressure [30]. The onset of coherence is at about $T^* \sim 20$ K, and no sign of superconductivity is found down to 0.1 K. Furthermore, 1D spin-chain behavior is also clear from susceptibility data [30], and density functional computations predict flat Fermi surfaces originating from the 1D f -electron bands along the c axis. An NFL state develops in a wide temperature range, as is apparent from the pressure-dependence resistivity data [30]. All these facts firmly suggest that CeCo_2Ga_8 is naturally positioned in the proximity of a magnetic QCP [30]. Nevertheless, T -linear resistivity and a logarithmically divergent specific heat are expected in a 2D antiferromagnetic QCP from the conventional Hertz-Millis theory [31,32], but not in a quasi-1D Kondo lattice system. Interestingly in the case of CeCo_2Ga_8 , anisotropic magnetic susceptibility behavior can be explained utilizing crystal-field theory, and the ratio of the exchange interaction is $|J_{\text{ex}}^c/J_{\text{ex}}^{a,b}| \sim 4\text{--}5$ [33]. Our investigation of CeCo_2Ga_8 includes electrical resistivity $\rho(T)$, dc susceptibility $\chi(T)$, and heat capacity $C_p(T)$ data used for characterization of CeCo_2Ga_8 and muon spin relaxation (μSR) measurement to study the low-energy spin dynamics. Our microscopic examination confirms that the stoichiometric CeCo_2Ga_8 compound exhibits an NFL ground state without any doping. It is interesting to note that an NFL ground state without doping is rare in Ce-based compounds, with only a few, CeRhBi [28,29], CeRhSn [34], and CeInPt_4 [35,36], belonging to this group.

II. EXPERIMENTAL DETAILS

For the present investigation, high-quality single crystals of CeCo_2Ga_8 were grown employing the Ga-flux method. The complete growth method is reported in Ref. [30]. The temperature-dependent magnetic susceptibility [$\chi(T)$] was measured using a Quantum Design Magnetic Property Measurement System (MPMS-SQUID) with the applied field parallel to the c axis. Electrical resistivity [$\rho(T)$] and heat capacity [$C_p(T)$] measurements were done using a Physical Property Measurement System (PPMS) with a dilution inset and He^3 cryostat, respectively.

μSR experiments were carried out the MUSR spectrometer in the zero-field (ZF) and longitudinal-field (LF; $H \parallel$ muon spin direction) geometries at the muon beamline of the ISIS Facility at the Rutherford Appleton Laboratory, Oxfordshire, U.K. [37]. Unaligned single crystals [as they were very small: less than 1.5 mm (length) \times 1 mm (diameter)] of CeCo_2Ga_8 were mounted on a 99.995% silver plate, applying thinned GE varnish covered with a high-purity silver foil. The sample was cooled down to 70 mK using a dilution refrigerator. The spin-polarized incident muon thermalized on the sample, and the resultant asymmetry was determined employing $G_z(t) = [N_F(t) - \alpha N_B(t)]/[N_F(t) + \alpha N_B(t)]$, where $N_B(t)$ and $N_F(t)$ are the number of positrons counted in the backward and forward detectors, respectively, and α is an instrumental calibration constant determined with a small (2-mT) transverse magnetic field at a high temperature. $G_z(t)$ gives information

about the spin-lattice relaxation rate and internal field distribution. We have collected ZF μSR data between 0.07 and 4 K, and LF μSR data in an applied field up to 0.3 T at 0.25 K. ZF/LF- μSR data were analyzed utilizing WiMDA software [38].

III. RESULTS AND DISCUSSION

Figure 1(a) represents the orthorhombic structure (space group $Pbam$, No. 55) of CeCo_2Ga_8 , showing the quasi-1D chains of Ce atoms along the c axis. The individual unit cell holds four Ce atoms. The NFL state in CeCo_2Ga_8 is confirmed by electrical resistivity, magnetization, and heat capacity measurements, and the results are presented in Figs. 1(b)–1(d). The temperature variation of susceptibility measured under a field-cooled condition with a 10 mT applied field parallel to the c axis is illustrated in Fig. 1(b). $\chi(T)$ manifests an $\sim T^{-0.2}$ response, suggesting CeCo_2Ga_8 located near the quantum phase transition [30]. This type of power-law behavior of $\chi(T)$ is also observed for other NFL systems such as CeRhBi [8,29]. On the other hand, the susceptibility of the NFL system $\text{CeCu}_{5.9}\text{Au}_{0.1}$ exhibits $\chi(T) \sim \chi_0(1 - a * T^{1/2})$ behavior [1]. The high-temperature Curie-Weiss fit yields an effective magnetic moment $\mu_{\text{eff}} = 2.74\mu_B$, which is larger compared to the free Ce^{3+} -ion value ($2.54\mu_B$); this may indicate a very weak magnetic contribution from the Co ion in CeCo_2Ga_8 . This peculiarity is well known in Ce-based HF systems, for instance, CeCoAsO [39] and CeCo_2As_2 [40]. In the low-temperature limit as shown in Fig. 1(b), $0.1 \text{ K} \leq T \leq 2 \text{ K}$, $\rho(T)$ varies linearly with the temperature ($\rho \sim T$), characteristic of an NFL state. The heat capacity varies logarithmically, $C_p/T \sim -\ln(T)$, in the low- T limit as shown in Fig. 1(d). It is clear from the inset in Fig. 1(d) that the heat capacity exhibits a broad maximum near 90 K, which can be attributed to the Schottky anomaly arising from the crystal-field effect in the presence of the Kondo effect. All of these attributes of CeCo_2Ga_8 are rather similar to the quasi-1D NFL ground state and lead to further examination using a microscopic technique such as muon spin relaxation measurement.

To probe the NFL state as seen from electrical, thermal, and magnetic measurements at low T , we carried out ZF/LF- μSR measurements [41]. The ZF depolarization reveals the sum of the local responses of muons embedded at different stopping sites in CeCo_2Ga_8 . ZF- μSR muon asymmetry spectra of CeCo_2Ga_8 at $T = 70$ mK (black symbols) and $T = 4$ K (blue symbols), which are representative of the data collected, are shown in Fig. 2(a). Both the 4 K and the 70 mK data in Fig. 2(a) reveal the same value of the initial asymmetry at $t = 0$ along with the lack of oscillations confirming the absence of long-range magnetic ordering in CeCo_2Ga_8 down to 70 mK. This precludes any argument for static magnetism in the sample [42]. Hence the moderate increase in the relaxation upon cooling from high temperatures reflects only a slowing-down of the electronic spin dynamics. Fits to the ZF- μSR data at different temperatures were done employing a Gaussian Kubo-Toyabe function multiplied by an exponential decaying function [43–45],

$$P_z(t) = A_1 \left[\frac{1}{3} + \frac{2}{3} (1 - \sigma_{\text{KT}}^2 t^2) \exp\left(\frac{-\sigma_{\text{KT}}^2 t^2}{2}\right) \right] \exp(-\lambda_{\text{ZF}} t) + A_{\text{bg}}, \quad (1)$$




FIG. 1. (a) Crystal structure of CeCo_2Ga_8 , showing the quasi-1D chains of Ce atoms along the c axis. The individual unit cell holds four Ce atoms. (b) Magnetic susceptibility as a function of temperature (log-log scale) parallel to the c axis ($\chi \sim T^{-0.2}$). (c) Low-temperature region of resistivity ($\rho \sim T$). Inset: $\rho(T)$ vs T plotted on the x axis as logarithmic scale down to 0.1 K. (d) Low-temperature dependence of heat capacity divided by temperature C_p/T plotted at a semilogarithmic scale in zero applied field ($C_p/T \sim -\ln T$). Inset: C_p/T over a large temperature range.

where λ_{ZF} is the ZF relaxation rate arising due to the local moment, and A_1 and A_{bg} are the asymmetries originating from the sample and background, respectively. A_{bg} was determined from the high-temperature ZF, which was kept fixed for the analysis. σ_{KT} is the nuclear contribution that emerges from the Gaussian distribution of the magnetic field at the muon site. The relaxation term $\exp(-\lambda_{\text{ZF}}t)$ is the magnetic contribution that comes from the fluctuating electronic spins, which provides information on the low-energy spin dynamics of CeCo_2Ga_8 . Considering that we have observed only a single-exponential function in ZF- μ SR data, this indicates that magnetic disorder in CeCo_2Ga_8 is negligible and hence the observed NFL behavior is intrinsic and has its origin in stoichiometric crystallographic-ordered CeCo_2Ga_8 .

As shown in the left panel in Fig. 2(b), the T variation of λ_{ZF} increases sharply below 1 K, indicating the development of a NFL state as evidenced by the bulk properties. Above 1 K, λ_{ZF} decreases with increasing temperature. The right panel in Fig. 2(b) represents the Arrhenius-like behavior of $\lambda_{\text{ZF}}(T)$, i.e., follows the form $\lambda_{\text{ZF}} = \lambda_0 \exp(-\frac{E_a}{k_B T})$, where E_a and k_B are the activation energy and Boltzmann constant, respectively. This confirms that the low- T spin dynamics of

CeCo_2Ga_8 is thermally activated with $E_a = 2.3$ mK, which is similar to observations of CeInPt_4 [36] and CeRhBi [29], with E_a values of 2.9 and 140 mK, respectively. It is noteworthy that the amplitude of the thermal fluctuations decreases with decreasing temperature, and at 70 mK thermal fluctuations are not important. We therefore attribute the observed behavior of the temperature dependence of $\lambda_{\text{ZF}}(T)$ to the quantum fluctuations, which is in agreement with the observed behavior of the heat capacity, resistivity, and magnetic susceptibility of CeCo_2Ga_8 . It is an open question why the temperature dependent relaxation of stoichiometric CeRhBi [29], CeInPt_4 [36] and CeCo_2Ga_8 compounds exhibits Arrhenius behavior in the NFL state (as $T \rightarrow 0$), while that of chemically disordered NFL systems such as $\text{CeRh}_{0.85}\text{Pd}_{0.15}$ exhibits power-law behavior [46]. We also plotted $\lambda_{\text{ZF}}(T)$ data for CeCo_2Ga_8 in a log-log plot [inset in Fig. 2(b)] to see the power-law behavior, but the data did not follow power-law behavior.

A longitudinal field of just 40 mT removes any relaxation due to a spontaneous field and is adequate to decouple muons from the relaxation channel, as presented in Fig. 2(c) at 250 mK. Once a muon is decoupled from the nuclear moments, the spectra can be best fitted using [47] $G_{z(t)} =$




FIG. 2. (a) Time-dependent zero-field μ SR spectra of CeCo_2Ga_8 collected at 70 mK (black symbols) and 4 K (blue symbols). Solid lines are the least-squares fit applying Eq. (1). (b) The left y axis plots the electronic contribution of the muon relaxation rate λ_{ZF} . A clear signature of quantum fluctuations is seen below 1 K, confirming the NFL state as shown by the bulk properties. The right axis demonstrates the Arrhenius behavior of λ_{ZF} . The line is the least-squares fit of the data, as presented in the text. Inset: A log-log plot of λ_{ZF} vs temperature. (c) Time-dependent longitudinal-field muon asymmetry spectra of CeCo_2Ga_8 measured at $T = 0.25$ K in zero field and at 40 mT. Solid lines are the least-squares fit to the raw data using Eq. (1). (d) Longitudinal-field dependence of the muon relaxation rate λ for CeCo_2Ga_8 at 0.25 K. The solid red line is the least-squares fit using Eq. (2).

$A_1 \exp(-\lambda_{\text{LF}}t) + A_{\text{bg}}$. λ_{LF} decreases rapidly at a low H and saturates at a high H . $\lambda_{\text{LF}}(H)$ can be adequately expressed by the standard description given by the Redfield formula [48,49],

$$\lambda = \lambda_0 + \frac{2\gamma_\mu^2 \langle H_l^2 \rangle \tau_c}{1 + \gamma_\mu^2 H^2 \tau_c^2}, \quad (2)$$

where λ_0 is the field-independent depolarization rate, and $\langle H_l^2 \rangle$ is the time-varying local field at muon sites due to the fluctuations of Ce $4f$ moments. H_l is the applied longitudinal field and the correlation time τ_c is related to the imaginary component of the q -independent dynamical susceptibility, $\chi''(w)$, through the fluctuation-dissipation theorem [50]. The red line in Fig. 2(d) represents the fit to the $\lambda_{\text{LF}}(H)$ data. The calculated parameters are $\lambda_0 = 0.19(1) \mu\text{s}^{-1}$, $\langle H_l^2 \rangle = 1.3(1) \text{ mT}$, and $\tau_c = 3.1(6) \times 10^{-8} \text{ s}$. The value of the time constant of CeCo_2Ga_8 unveils a slow spin dynamics, which originates from the quantum critical fluctuations at low T . Similar values are observed for CeRhBi [29]: $\lambda_0 = 0.17(1) \mu\text{s}^{-1}$, $\langle H_l^2 \rangle = 1.5(1) \text{ mT}$, and $\tau_c = 4.2(6) \times 10^{-8} \text{ s}$.

IV. CONCLUSION

In conclusion, we have presented magnetization, resistivity, heat capacity, and ZF/LF muon spin relaxation measurements on the quasi-one-dimensional CeCo_2Ga_8 compound. The linear behavior of $\rho(T)$, power-law divergence of $\chi(T) \sim T^{-0.2}$, and logarithmic divergence of $C_p(T)/T$ suggest a NFL ground state of CeCo_2Ga_8 . Moreover, the increase in the ZF relaxation rate λ_{ZF} below 1 K and the Arrhenius-like behavior of CeCo_2Ga_8 suggest a NFL ground state, which is quite similar to that seen in other NFL systems, for example, CeRhBi [29] and CeInPt_4 [36]. Our ZF μ SR measurements confirm the absence of long-range magnetic ordering down to 70 mK. Furthermore, the longitudinal-field dependence μ SR study provides information on the spin fluctuation rate and the width of the field distribution at the muon sites. The observed quantum fluctuations below 1 K in the undoped CeCo_2Ga_8 compound make it a prototype material for investigation of the low- T quantum fluctuations in low-dimensional NFL systems and other Ce-128 counterparts with a YbCo_2Ga_8 -type structure. This work will pave the way to our understanding of the NFL in 1D systems, both theoretically and experimentally.

ACKNOWLEDGMENTS

We would like to thank Dr. J. Sannigrahi and Dr. V. K. Anand for interesting discussions. A.B. would like to thank the Department of Science and Technology (DST), India, for an Inspire Faculty Research Grant (DST/INSPIRE/04/2015/000169), the DST, India (SR/NM/Z-07/2015), for financial support, and the Jawaharlal Nehru Centre for Advanced Scientific Research (JNCASR) for management of the project. D.T.A. is grateful to the Royal Society of London (RSL) for the U.K.-China Newton mobility grant, and D.T.A. and H.L. acknowledge the funding support of the Newton Advanced Fellowship from the RSL (Grant No. NAF_R1_201248). K.P. would like to acknowledge the DST, India, for an Inspire Fellowship (No. IF170620). Work

at the Institute of Physics, Chinese Academy of Sciences (CAS), was supported by the National Key Research and Development Program of China (Grants No. 2017YFA0303100 and No. 2017YFA0302900), the National Natural Science Foundation of China (Grants No. 11974392, No. 11822411, No. 11774399, No. 11961160699, and No. 12061130200), and the Beijing Natural Science Foundation (Grants No. JQ19002 and No. Z180008). H.L. is grateful for the support from the Youth Innovation Promotion Association of the CAS (Grant No. 2016004). A.M.S. thanks South African National Research Foundation (SA-NRF) and the University Research Committee/Faculty Research Committee of the University of Johannesburg for generous financial assistance. We would like to thank the ISIS Facility for beam time (Experiment no: RB1810451) [37].

- [1] H. v. Löhneysen, A. Rosch, M. Vojta, and P. Wölfle, *Rev. Mod. Phys.* **79**, 1015 (2007).
- [2] T. Shibauchi, A. Carrington, and Y. Matsuda, *Annu. Rev. Condens. Matter Phys.* **5**, 113 (2014).
- [3] I. Fischer and A. Rosch, *Phys. Rev. B* **71**, 184429 (2005).
- [4] A. Georges, G. Kotliar, W. Krauth, and M. J. Rozenberg, *Rev. Mod. Phys.* **68**, 13 (1996).
- [5] C. M. Varma, *Rev. Mod. Phys.* **48**, 219 (1976).
- [6] P. Coleman, in *Handbook of Magnetism and Advanced Magnetic Materials*, edited by H. Knoemuller and S. Parkin (Wiley, New York, 2007), Vol. 1, p. 95.
- [7] P. S. Riseborough, *Adv. Phys.* **49**, 257 (2000); *Phys. Rev. B* **45**, 13984 (1992).
- [8] G. R. Stewart, *Rev. Mod. Phys.* **73**, 797 (2001); A. Amato, *ibid.* **69**, 1119 (1997).
- [9] L. D. Landau, *J. Expt. Theoret. Phys. (USSR)* **30**, 1058 (1956) [*Sov. Phys. JETP* **3**, 920 (1957)].
- [10] G. Baym and C. Pethick, *Landau Fermi-Liquid Theory: Concepts and Applications* (John Wiley & Sons, New York, 2008).
- [11] C. Krellner, S. Lausberg, A. Steppke, M. Brando, L. Pedrero, H. Pfau, S. Tencé, H. Rosner, F. Steglich, and C. Geibel, *New J. Phys.* **13**, 103014 (2011).
- [12] A. Steppke, R. Küchler, S. Lausberg, E. Lengyel, L. Steinke, R. Borth, T. Lühmann, C. Krellner, M. Nicklas, C. Geibel, F. Steglich, and M. Brando, *Science* **339**, 933 (2013).
- [13] L. D. Landau, and E. M. Lifshitz, *Statistical Physics, Part 2* (Pergamon, Oxford, UK, 1980).
- [14] A. J. Schofield, *Contemp. Phys.* **40**, 95 (1999).
- [15] G. R. Stewart, *Rev. Mod. Phys.* **78**, 743 (2006).
- [16] P. Gegenwart, Q. Si, and F. Steglich, *Nat. Phys.* **4**, 186 (2008).
- [17] Q. Si and F. Steglich, *Science* **329**, 1161 (2010).
- [18] O. Stockert and F. Steglich, *Annu. Rev. Condens. Matter Phys.* **2**, 79 (2011).
- [19] U. Schollwöck, *Rev. Mod. Phys.* **77**, 259 (2005).
- [20] H. von Löhneysen, *J. Phys.: Condens. Matter* **8**, 9689 (1996).
- [21] K. Heuser, E.-W. Scheidt, T. Schreiner, and G. R. Stewart, *Phys. Rev. B* **57**, R4198 (1998).
- [22] H. Shishido, T. Shibauchi, K. Yasu, T. Kato, H. Kontani, T. Terashima, and Y. Matsuda, *Science* **327**, 980 (2010).
- [23] L. P. Gor'kov and P. D. Grigoriev, *Phys. Rev. B* **73**, 060401(R) (2006).
- [24] N. D. Mathur, F. M. Grosche, S. R. Julian, I. R. Walker, D. M. Freye, R. K. W. Haselwimmer, and G. G. Lonzarich, *Nature* **394**, 39 (1998).
- [25] A. P. Ramirez, P. Chandra, P. Coleman, Z. Fisk, J. L. Smith, and H. R. Ott, *Phys. Rev. Lett.* **73**, 3018 (1994).
- [26] F. Bommeli, L. Degiorgi, P. Wachter, F. B. Anders, and A. V. Mitin, *Phys. Rev. B* **56**, R10001(R) (1997).
- [27] F. Steglich, P. Hellmann, S. Thomas, P. Gegenwart, A. Link, R. Helfrich, G. Sparn, M. Lang, C. Geibel, and W. Assmus, *Physica B: Condens. Matter* **237**, 192 (1997).
- [28] T. Sasakawa, K. Shigetoh, D. Hirata, K. Umeo, and T. Takabatake, *Physica B: Condens. Matter* **359–361**, 111 (2005).
- [29] V. K. Anand, D. T. Adroja, A. D. Hillier, K. Shigetoh, T. Takabatake, J. G. Park, K. A. McEwen, J. H. Pixley, and Q. Si, *J. Phys. Soc. Jpn.* **87**, 064708 (2018).
- [30] L. Wang, Z. Fu, J. Sun, M. Liu, W. Yi, C. Yi, Y. Luo, Y. Dai, G. Liu, Y. Matsuhita, K. Yamaura, L. Lu, J. G. Cheng, Y. F. Yang, Y. Shi, and J. Luo, *npj Quantum Mater.* **2**, 36 (2017).
- [31] J. A. Hertz, *Phys. Rev. B* **14**, 1165 (1976).
- [32] A. J. Millis, *Phys. Rev. B* **48**, 7183 (1993).
- [33] K. Cheng, L. Wang, Y. Xu, F. Yang, H. Zhu, J. Ke, X. Lu, Z. Xia, J. Wang, Y. Shi, Y. Yang, and Y. Luo, *Phys. Rev. Materials* **3**, 021402 (2019).
- [34] P. C. Ho, V. S. Zapf, A. Ślebarski, and M. B. Maple, *Philos. Mag.* **84**, 2119 (2004).
- [35] S. K. Malik, D. T. Adroja, M. Slaski, B. D. Dunlap, and A. Umezawa, *Phys. Rev. B* **40**, 9378(R) (1989).
- [36] A. D. Hillier, D. T. Adroja, S. R. Giblin, W. Kockelmann, B. D. Rainford, and S. K. Malik, *Phys. Rev. B* **76**, 174439 (2007).
- [37] D. Adroja, A. Bhattacharyya, and J. Lord, *Understand the Role of Dimensionality on Heavy Fermion Physics and Quantum Criticality in CeCo₂Ga₈: A Muon Study* (ISIS Neutron and Muon Source, Oxfordshire, UK, 2018).
- [38] F. L. Pratt, *Physica B* **289–290**, 710 (2000).
- [39] R. Sarkar, A. Jesche, C. Krellner, M. Baenitz, C. Geibel, C. Mazumdar, and A. Poddar, *Phys. Rev. B* **82**, 054423 (2010).

- [40] K. W. H. Stevens, *Proc. Phys. Soc., Sec. A* **65**, 209 (1952).
- [41] A. Bhattacharyya, D. T. Adroja, and M. Smidman, *Sci. China Phys. Mech. Astron.* **61**, 127402 (2018).
- [42] T. Suzuki, I. Watanabe, F. Yamada, Y. Ishii, K. Ohishi, Risdiana, T. Goto, and H. Tanaka, *Phys. Rev. B* **80**, 064407 (2009).
- [43] A. Bhattacharyya, C. Ritter, D. T. Adroja, F. C. Coomer, and A. M. Strydom, *Phys. Rev. B* **94**, 014418 (2016).
- [44] A. Bhattacharyya, D. T. Adroja, K. Panda, S. Saha, T. Das, A. J. S. Machado, O. V. Cigarroa, T. W. Grant, Z. Fisk, A. D. Hillier, and P. Manfrinetti, *Phys. Rev. Lett.* **122**, 147001 (2019).
- [45] A. Bhattacharyya, P. P. Ferreira, F. B. Santos, D. T. Adroja, J. S. Lord, L. E. Correa, A. J. S. Machado, A. L. R. Manesco, and L. T. F. Eleno, *Phys. Rev. Res.* **2**, 022001(R) (2020).
- [46] D. T. Adroja, A. D. Hillier, J.-G. Park, W. Kockelmann, K. A. McEwen, B. D. Rainford, K.-H. Jang, C. Geibel, and T. Takabatake, *Phys. Rev. B* **78**, 014412 (2008).
- [47] R. S. Hayano, Y. J. Uemura, J. Imazato, N. Nishida, T. Yamazaki, and R. Kubo, *Phys. Rev. B* **20**, 850 (1979).
- [48] A. Yaouanc and P. Dalmas De Reotier, *Muon Spin Rotation, Relaxation, and Resonance: Applications to Condensed Matter* (Oxford University Press, Oxford, UK, 2011).
- [49] A. Bhattacharyya, D. T. Adroja, A. M. Strydom, A. D. Hillier, J. W. Taylor, A. Thamizhavel, S. K. Dhar, W. A. Kockelmann, and B. D. Rainford, *Phys. Rev. B* **90**, 054405 (2014).
- [50] J. S. Toll, *Phys. Rev.* **104**, 1760 (1956).