



# On the oracle complexity of first-order and derivative-free algorithms for smooth nonconvex minimization

C Cartis, NIM Gould, PL Toint

October 2010

**©2010 Science and Technology Facilities Council**

Enquiries about copyright, reproduction and requests for additional copies of this report should be addressed to:

RAL Library  
Science and Technology Facilities Council  
Rutherford Appleton Laboratory  
Harwell Science and Innovation Campus  
Didcot  
OX11 0QX

Tel: +44(0)1235 445384  
Fax: +44(0)1235 446403  
email: [library@rl.ac.uk](mailto:library@rl.ac.uk)

Science and Technology Facilities Council reports are available online at: <http://epubs.cclrc.ac.uk/>

**ISSN 1358-6254**

Neither the Council nor the Laboratory accept any responsibility for loss or damage arising from the use of information contained in any of their reports or in any communication about their tests or investigations.

# On the oracle complexity of first-order and derivative-free algorithms for smooth nonconvex minimization

Coralia Cartis<sup>1</sup>, Nicholas I. M. Gould<sup>2,3</sup> and Philippe L. Toint<sup>4</sup>

## ABSTRACT

The (optimal) function/gradient evaluations worst-case complexity analysis available for the Adaptive Regularizations algorithms with Cubics (ARC) for nonconvex smooth unconstrained optimization is extended to finite-difference versions of this algorithm, yielding complexity bounds for first-order and derivative free methods applied on the same problem class. A comparison with the results obtained for derivative-free methods by Vicente (2010) is also discussed, giving some theoretical insight on the relative merits of various methods in this popular class of algorithms.

---

<sup>1</sup> School of Mathematics, The King's Buildings, University of Edinburgh, EH9 3JZ, Scotland, EU. Email: coralia.cartis@ed.ac.uk .

<sup>2</sup> Computational Science and Engineering Department, Rutherford Appleton Laboratory, Chilton, Oxfordshire, OX11 0QX, England, EU. Email: nick.gould@stfc.ac.uk .  
Current reports available from "<http://www.numerical.rl.ac.uk/reports/reports.shtml>".

<sup>3</sup> This work was supported by the EPSRC grant EP/E053351/1.

<sup>4</sup> Department of Mathematics, Facultés Universitaires ND de la Paix, 61, rue de Bruxelles, B-5000 Namur, Belgium, EU. Email : philippe.toint@fundp.ac.be .  
Current reports available from "<http://www.fundp.ac.be/~phtoint/pht/publications.html>".

Computational Science and Engineering Department  
Atlas Centre  
Rutherford Appleton Laboratory  
Oxfordshire OX11 0QX

October 21, 2010

# 1 Introduction

We consider algorithms for the solution of the unconstrained (possibly nonconvex) optimization problem

$$\min_x f(x) \tag{1.1}$$

where we assume that  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is smooth (in a sense to be specified later) and bounded below. All methods for the solution of (1.1) are iterative and, starting from some initial guess  $x_0$ , generate a sequence  $\{x_k\}$  of iterates approximating a critical point of  $f$ . A variety of algorithms of this form exist, and they are often classified according to their requirements in terms of computing derivatives of the objective function. First-order methods are methods which use  $f(x)$  and its gradient  $\nabla_x f(x)$ , and derivative-free (or zeroth order) methods are those which only use  $f(x)$ , without any gradient computation. This paper is concerned with estimating worst-case bounds on the number of objective function and/or gradient calls that are necessary for the specific methods in these two classes to compute approximate critical points for (1.1), starting from arbitrary initial guesses  $x_0$ . These bound in turn provide upper bounds on the complexity of solving (1.1) with general algorithms in the first-order or derivative-free classes.

Worst-case complexity analysis for optimization methods probably really started with Nemirovski and Yudin (1983), where the notion of oracle (or black-box) complexity was introduced. Instead of expressing complexity in terms of simple operation counts, the complexity of an algorithm is measured by the number of calls this algorithm makes, in the worst-case, to an oracle (the computation of the objective function or the gradient values, for instance) in order to successfully terminate. Many results of that nature have been derived since, mostly on the convex optimization problem (see, for instance, Nesterov 2004, 2008, Nemirovski, 1994, Agarwal, Bartlett, Ravikummar and Wainwright, 2009), but also for the nonconvex case (see Vavasis 1992*b*, 1992*a*, 1993, Nesterov and Polyak, 2006, Gratton, Sartenaer and Toint, 2008, Cartis, Gould and Toint 2009*a*, 2010*a*, 2010*b*, 2010*c*, or Vicente, 2010). Of particular interest here is the Adaptive Regularizations with Cubics (ARC) algorithm independently proposed by Griewank (1981), Weiser et al (2007) and Nesterov and Polyak (2006), whose worst-case iteration complexity<sup>1</sup> was shown in the last of these references to be of  $O(\epsilon^{-3/2})$  for finding an approximate solution  $x_*$  such that the gradient at  $x_*$  is smaller than  $\epsilon$  in norm. This result was extended by Cartis et al. (2010*a*) to an algorithm no longer requiring the computation of exact second-derivatives, but merely of a suitably accurate approximation<sup>2</sup>. Moreover, Cartis et al. (2010*b*, 2010*c*) showed that, when exact second derivatives are used, this complexity bound is tight and is optimal within a large class of second-order methods.

The purpose of the present paper is to use the freedom left in Cartis et al. (2010*a*)

---

<sup>1</sup>That is its oracle complexity for a choice of the oracle corresponding to the computation of the objective function and its first and second derivatives.

<sup>2</sup>This method also abandoned global optimization of the underlying cubic model and avoided an *a priori* knowledge of the objective function's Hessian's Lipschitz constant, two assumptions made by Nesterov and Polyak (2006).

to approximate the objective function's Hessian to explore complexity bounds for finite-difference methods in exact arithmetic, and thereby establish lower bounds on the oracle complexity of methods for solving unconstrained nonconvex problems, where the oracle consists of evaluating objective-function and/or gradient values. The ARC algorithm and the associated complexity bounds are recalled in Section 2. Section 3 investigates the case of a first-order variant in which the objective-function's Hessian is approximated by finite differences in gradient values, while Section 4 considers a derivative-free variant where the gradient of  $f$  is computed by central differences and its Hessian by forward differences. These results are finally discussed and compared to existing complexity bounds by Vicente (2010) in Section 5.

## 2 The ARC algorithm and its oracle complexity

The Adaptive Regularization with Cubics (ARC) algorithm is based on the approximate minimization, at iteration  $k$ , of (the possibly nonconvex) cubic model

$$m_k(s) = \langle g_k, s \rangle + \frac{1}{2} \langle s, B_k s \rangle + \frac{1}{3} \sigma_k \|s\|^3, \quad (2.1)$$

where  $\langle \cdot, \cdot \rangle$  denotes the Euclidean inner product and  $\|\cdot\|$  the Euclidean norm. Here  $B_k$  is a symmetric  $n \times n$  approximation of  $\nabla_{xx} f(x_k)$ ,  $\sigma_k > 0$  is a regularization weight and

$$g_k = \nabla_x m_k(0) = \nabla_x f(x_k). \quad (2.2)$$

By ‘‘approximate minimization’’, we mean that a step  $s_k$  is computed that satisfies

$$\langle g_k, s_k \rangle + \langle s_k, B_k s_k \rangle + \sigma_k \|s_k\|^3 \leq 0, \quad (2.3)$$

$$\langle s_k, B_k s_k \rangle + \sigma_k \|s_k\|^3 \geq 0 \quad (2.4)$$

$$m_k(s_k) \leq m_k(s_k^C) \quad (2.5)$$

with

$$s_k^C = -\alpha_k^C g_k \quad \text{and} \quad \alpha_k^C = \arg \min_{\alpha \leq 0} m_k(-\alpha g_k), \quad (2.6)$$

and

$$\|\nabla_x m_k(s_k)\| = \|g_k + B_k s_k + \sigma_k \|s_k\| s_k\| \leq \kappa_\theta \min[1, \|s_k\|] \|g_k\|, \quad (2.7)$$

for some given constant  $\kappa_\theta \in (0, 1)$ .

As noted in Cartis et al. (2010a), conditions (2.3) and (2.4) must hold if  $s_k$  minimizes the model along the direction  $s_k/\|s_k\|$ , while (2.7) holds by continuity if  $s_k$  is sufficiently close to a first-order critical point of  $m_k$ . Moreover, (2.5)-(2.6) are nothing but the familiar Cauchy-point decrease condition. Fortunately, these conditions can be ensured algorithmically. In particular, conditions (2.3)–(2.7) hold if  $s_k$  is a (computable) global minimizer of  $m_k$  (see Griewank, 1981, Nesterov and Polyak, 2006, see also Cartis, Gould and Toint, 2009c). Note that, since  $\nabla_x m_k(0) = \nabla_x f(x_k)$ , (2.7) may be interpreted as requiring a relative reduction in the norm of the model's gradient at least equal to  $\kappa_\theta \min[1, \|s_k\|]$ .

The ARC algorithm may then be stated as presented on the facing page.

**Algorithm 2.1: ARC**

**Step 0:** An initial starting point  $x_0$  is given, as well as a user-defined accuracy threshold  $\epsilon \in (0, 1)$ . Set  $k = 0$ .

**Step 1:** If  $\|\nabla_x f(x_k)\| \leq \epsilon$ , terminate with approximate solution  $x_k$ .

**Step 2:** Compute any Hessian approximation  $B_k$ .

**Step 3:** Compute a step  $s_k$  satisfying (2.3)–(2.7).

**Step 4:** Compute  $f(x_k + s_k)$  and

$$\rho_k = \frac{f(x_k) - f(x_k + s_k)}{-m_k(s_k)}. \quad (2.8)$$

**Step 5:** Set

$$x_{k+1} = \begin{cases} x_k + s_k & \text{if } \rho_k \geq \eta_1, \\ x_k & \text{otherwise.} \end{cases}$$

**Step 6:** Set

$$\sigma_{k+1} \in \begin{cases} (0, \sigma_k] & \text{if } \rho_k > \eta_2, & \text{[very successful iteration]} \\ [\sigma_k, \gamma_1 \sigma_k] & \text{if } \eta_1 \leq \rho_k \leq \eta_2, & \text{[successful iteration]} \\ [\gamma_1 \sigma_k, \gamma_2 \sigma_k] & \text{otherwise.} & \text{[unsuccessful iteration]} \end{cases} \quad (2.9)$$

**Step 7:** Increment  $k$  by one and return to Step 1.

In this algorithm, we assume that the constants satisfy  $\gamma_2 \geq \gamma_1 > 1$ ,  $1 > \eta_2 \geq \eta_1 > 0$  and  $\sigma_0 > 0$ . We denote by

$$\mathcal{S} = \{k \geq 0 \mid \text{iteration } k \text{ is successful or very successful in the sense of (2.9)}\}$$

the set of *successful iterations*, and

$$\mathcal{S}_j = \{k \in \mathcal{S} \mid k \leq j\} \quad \text{and} \quad \mathcal{U}_j = \{0, \dots, j\} \setminus \mathcal{S}_j, \quad (2.10)$$

the sets of successful and unsuccessful iterations up to iteration  $j$ .

It is not the purpose of the present paper to discuss implementation issues or convergence theory for the ARC algorithm, but we need to recall the main complexity results for this method, as well as the assumptions under which these hold.

We first restate our assumptions.

**A.1:** The objective function  $f$  is twice continuously differentiable on  $\mathbb{R}^n$  and its gradient and Hessian are Lipschitz continuous on the path of iterates with Lipschitz constant

$L_g$  and  $L_H$ , respectively, i.e., for all  $k \geq 0$  and all  $\alpha \in [0, 1]$ ,

$$\|\nabla_x f(x_k) - \nabla_x f(x_k + \alpha s_k)\| \leq L_g \alpha \|s_k\| \quad (2.11)$$

and

$$\|\nabla_{xx} f(x_k) - \nabla_{xx} f(x_k + \alpha s_k)\| \leq L_H \alpha \|s_k\|. \quad (2.12)$$

**A.2:** The objective function  $f$  is bounded below, that is there exists a constant  $f_{\text{low}}$  such that

$$f(x) \geq f_{\text{low}}$$

for all  $x \in \mathbb{R}^n$

**A.3:** For all  $k \geq 0$ , the Hessian approximation  $B_k$  satisfies

$$\|B_k\| \leq \kappa_B \quad (2.13)$$

and

$$\|(\nabla_{xx} f(x_k) - B_k)s_k\| \leq \kappa_{\text{BH}} \|s_k\|^2 \quad (2.14)$$

for some constants  $\kappa_B > 0$  and  $\kappa_{\text{BH}} > 0$ .

We start by noting that the form of the cubic model (2.1) ensures a remarkable bound on the the stepnorm and model decrease.

**Lemma 2.1.** *Suppose that (2.3), (2.4) and (2.5) hold. Then*

$$\|s_k\| \leq \frac{3}{\sigma_k} \max \left[ \|B_k\|, \sqrt{\sigma_k \|g_k\|} \right] \quad (2.15)$$

and

$$m_k(s_k) \leq -\frac{1}{6} \sigma_k \|s_k\|^3. \quad (2.16)$$

**Proof.** See Lemma 2.2 in Cartis et al. (2009a) for the proof of (2.15) and Lemma 4.2 in Cartis, Gould and Toint (2009b) for that of (2.16).  $\square$

For our purposes it is also useful to consider the following bounds on the value of the regularization parameter.

**Lemma 2.2.** *Suppose that A.1 and that (2.13) hold. Then there exists a constant  $\kappa_\sigma > 0$  independent of  $n$  such that, for all  $k \geq 0$*

$$\sigma_k \leq \max \left[ \sigma_0, \frac{\kappa_\sigma}{\epsilon} \right]. \quad (2.17)$$

*If, in addition, (2.14) also holds, then there exists a constant  $\sigma_{\text{max}} > 0$  independent of  $n$  and  $\epsilon$  such that, for all  $k \geq 0$*

$$\sigma_k \leq \sigma_{\text{max}}. \quad (2.18)$$

**Proof.** See Lemmas 3.2 and 3.3 in Cartis et al. (2010a) for the proof of (2.17) and Lemma 5.2 in Cartis et al. (2009a) for that of (2.18). Note that both of these proofs crucially depends on the identity (2.2), which means they have to be revisited if this equality fails.  $\square$

Without loss of generality, we assume in what follows that  $\epsilon$  is small enough for the second term in the max of (2.17) to dominate, and thus that (2.17) may be rewritten to state that, for all  $k \geq 0$

$$\sigma_k \leq \frac{\kappa_\sigma}{\epsilon}. \quad (2.19)$$

If (2.18) holds, then, crucially, the step  $s_k$  can then be proved to be sufficiently long compared to the gradient's norm at iteration  $k + 1$ .

**Lemma 2.3.** *Suppose that **A.1** and **A.3** hold. Then, for all  $k \geq 0$ , one has that, for some  $\kappa_g > 0$  independent of  $n$ ,*

$$\|s_k\| \geq \kappa_g \sqrt{\|\nabla_x f(x_k + s_k)\|}. \quad (2.20)$$

**Proof.** See Lemma 5.2 in Cartis et al. (2010a).  $\square$

The final important observation in the complexity analysis is that the total number of iterations required by the ARC algorithm to terminate may be bounded in terms of the number of successful iterations needed.

**Lemma 2.4.** *Suppose that **A.1** and **A.3** hold and, for any fixed  $j \geq 0$ , let  $\mathcal{S}_j$  and  $\mathcal{U}_j$  be defined in (2.10). Assume also that*

$$\sigma \geq \sigma_{\min} \quad (2.21)$$

for some  $\sigma_{\min} > 0$ . Then one has that

$$|\mathcal{U}_j| \leq \left[ (|\mathcal{S}_j| + 1) \frac{1}{\log \gamma_1} \log \left( \frac{\sigma_{\max}}{\sigma_{\min}} \right) \right]. \quad (2.22)$$

**Proof.** See Theorem 2.1 in Cartis et al. (2010a). Observe that this proof uniquely depends on the mechanism used in the algorithm for updating  $\sigma_k$ , and its independent of the values of  $g_k$  or  $B_k$ .  $\square$

Combining those results and using **A.2** then yields the following oracle complexity theorem.

**Theorem 2.5.** *Suppose that **A.1–A.3** hold, that  $\epsilon \in (0, 1)$  is given and that (2.21) holds. Then the ARC algorithm terminates after at most*

$$N_1^s \stackrel{\text{def}}{=} 1 + \lceil \kappa_S^s \epsilon^{-3/2} \rceil, \quad (2.23)$$

successful iterations and at most

$$N_1 \stackrel{\text{def}}{=} \lceil \kappa_S \epsilon^{-3/2} \rceil \quad (2.24)$$



iterations in total, where

$$\kappa_S^s \stackrel{\text{def}}{=} (f(x_0) - f_{\text{low}})/(\eta_1 \alpha_S), \quad \alpha_S \stackrel{\text{def}}{=} (\sigma_{\min} \kappa_g^3)/6 \quad (2.25)$$

and

$$\kappa_S \stackrel{\text{def}}{=} (1 + \kappa_S^u)(2 + \kappa_S^s), \quad \kappa_S^u \stackrel{\text{def}}{=} \log(\sigma_{\max}/\sigma_{\min})/\log \gamma_1, \quad (2.26)$$

with  $\kappa_g$  and  $\bar{\sigma}$  defined in (2.20) and (2.18), respectively. As a consequence, the ARC algorithm terminates after at most  $N_1^s$  gradient evaluations and at most  $N_1$  objective function evaluations.

**Proof.** See Corollary 5.3 in Cartis et al. (2010a). □

The bound given by (2.23) is known to be qualitatively<sup>3</sup>tight and optimal for a wide class of second-order methods (see Cartis et al. 2010b, 2010c). Also note that the constants in (2.25) and (2.26) do not depend on  $n$ .

### 3 A first-order finite-difference ARC variant

The objective of this section is to extend the ARC algorithm to a version using finite differences in gradients to compute the Hessian approximation  $B_k$ . If the accuracy of the finite-difference scheme is high enough to ensure that (2.14) holds, then one might expect that a worst-case iteration complexity similar to (2.23)-(2.24) would hold, thereby providing a first worst-case oracle complexity estimate for first-order methods applied on nonconvex unconstrained problems.

For defining this algorithm, which we will refer to as Algorithm ARC-FDH, we only need to specify the details of the estimation of  $B_k$ . If we compute this latter matrix by using  $n$  forward gradient differences at  $x_k$  with stepsize  $h_k$  of the form

$$\frac{\nabla_x f(x_k) - \nabla_x f(x_k + h_k e_j)}{h_k} \quad (3.1)$$

(where  $e_j$  is the  $j$ -th vector of the canonical basis) and symmetrize the result, it is well known (see Nocedal and Wright, 1999, Section 7.1) that

$$\|\nabla_{xx} f(x_k) - B_k\| \leq \kappa_{\text{eHg}} h_k \quad (3.2)$$

for some constant  $\kappa_{\text{eHg}} \in [0, L_H]$ . The only remaining issue is therefore to define a procedure guaranteeing that

$$h_k \leq \kappa_{\text{hs}} \|s_k\|. \quad (3.3)$$

for some  $\kappa_{\text{hs}} > 0$  and all  $k \geq 0$ . As we show below, this can be achieved if we consider Algorithm ARC-FDH on the next page, where  $\gamma_3 \in (0, 1)$  and  $\kappa_{\text{hs}} \geq 1$ .

---

<sup>3</sup>The constants may not be optimal.

**Algorithm 3.1: ARC-FDH**

**Step 0:** An initial starting point  $x_0$  is given, as well as a user-defined accuracy threshold  $\epsilon \in (0, 1)$ . If  $\|\nabla_x f(x_0)\| \leq \epsilon$ , terminate. Otherwise, set  $k = 0$ ,  $j = 0$  and choose an initial stepsize  $h_{0,0} \in (0, 1]$ .

**Step 1:** Estimate  $B_{k,j}$  using  $n$  gradient differences of the form (3.1), using the stepsize  $h_{k,j}$ .

**Step 2:** Compute a step  $s_{k,j}$  satisfying (2.3)–(2.7).

**Step 3:** Compute  $\nabla_x f(x_k + s_{k,j})$ . If  $\|\nabla_x f(x_k + s_{k,j})\| \leq \epsilon$ , terminate with approximate solution  $x_k + s_{k,j}$ .

**Step 4:** If

$$h_{k,j} > \kappa_{\text{hs}} \|s_{k,j}\|, \quad (3.4)$$

set  $h_{k,j+1} = \gamma_3 h_{k,j}$ , increment  $j$  by one and return to Step 1. Otherwise, set  $s_k = s_{k,j}$  and  $h_k = h_{k,j}$ .

**Step 5:** Compute  $f(x_k + s_k)$  and

$$\rho_k = \frac{f(x_k) - f(x_k + s_k)}{-m_k(s_k)}. \quad (3.5)$$

**Step 6:** Set

$$x_{k+1} = \begin{cases} x_k + s_k & \text{if } \rho_k \geq \eta_1, \\ x_k & \text{otherwise.} \end{cases}$$

**Step 7:** Set

$$\sigma_{k+1} \in \begin{cases} (0, \sigma_k] & \text{if } \rho_k > \eta_2, & \text{[very successful iteration]} \\ [\sigma_k, \gamma_1 \sigma_k] & \text{if } \eta_1 \leq \rho_k \leq \eta_2, & \text{[successful iteration]} \\ [\gamma_1 \sigma_k, \gamma_2 \sigma_k] & \text{otherwise.} & \text{[unsuccessful iteration]} \end{cases} \quad (3.6)$$

**Step 8:** Set  $h_{k+1,0} = h_k$  and  $j = 0$ . Increment  $k$  by one and return to Step 1 if  $\rho_k \geq \eta_1$ , or to Step 2 otherwise.

By convention and analogously to our notation for  $s_k$  and  $h_k$ , we denote by  $B_k$  the approximation  $B_{k,j}$  obtained at the end of the loop between Steps 1 and 4. Clearly, the test (3.4) in Step 4 ensures that (3.3) holds, as requested. Observe that, because the norm of the step is a monotonically decreasing function as a function of  $\sigma_k$  (see Lemma 3.1 in Cartis et al., 2009c), it decreases at an unsuccessful iteration, which might then possibly

require a new evaluation of the approximate Hessian in order to preserve (3.3). Observe also that the mechanism of the algorithm implies that the positive sequence  $\{h_k\}$  is monotonically decreasing and bounded above by  $h_{0,0} \leq 1$ .

It now remains to show that this algorithm is well defined, which we do under the additional assumption that the (true) gradients remain bounded at all iterates. Since the sequence  $\{f(x_k)\}$  is monotonically decreasing, this condition can for instance be ensured by assuming bounded gradients of the level set  $\{x \in \mathbb{R}^n \mid f(x) \leq f(x_0)\}$ .

**A.4:** There exists a constant  $\kappa_{\text{ubg}} \geq 0$  such that, for all  $k \geq 0$

$$\|\nabla_x f(x_k)\| \leq \kappa_{\text{ubg}}.$$

**Lemma 3.1.** *Suppose that **A.1**, **A.4** and (2.21) hold, Then (2.13) holds with*

$$\kappa_B \stackrel{\text{def}}{=} \max[\kappa_{\text{eHg}} + L_g, \sqrt{\kappa_\sigma \kappa_{\text{ubg}}}] \geq \sqrt{\kappa_\sigma \kappa_{\text{ubg}}}$$

and, for all  $k \geq 0$ ,

$$\|s_k\| \geq \frac{(1 - \kappa_\theta) \epsilon}{\max[4\kappa_B, \kappa_B + 3\sqrt{\sigma_k \kappa_{\text{ubg}}}]}. \quad (3.7)$$

**Proof.** We first note that (2.11) ensures that  $\|\nabla_{xx} f(x_k)\| \leq L_g$  for all  $k \geq 0$  and therefore that

$$\|B_{k,j}\| \leq \|B_{k,j} - \nabla_{xx} f(x_k)\| + \|\nabla_{xx} f(x_k)\| \leq \kappa_{\text{eHg}} + L_g \leq \max[\kappa_{\text{eHg}} + L_g, \sqrt{\kappa_\sigma \kappa_{\text{ubg}}}], \quad (3.8)$$

where we used the triangle inequality, the bound  $h_{k,j} \leq h_{0,0} \leq 1$  and (3.2). Hence (2.13) holds. Observe now that (2.2) and the mechanism of the algorithm then implies that, as long as the algorithm hasn't terminated,

$$\|g_k\| > \epsilon. \quad (3.9)$$

We know from (2.7) and (2.2) that, for all  $k \geq 0$ ,

$$\kappa_\theta \min[1, \|s_k\|] \|g_k\| \geq \|\nabla_x m_k(0) + B_k s_k + \sigma_k \|s_k\| s_k\| \geq \|g_k\| - \|B_k s_k + \sigma_k \|s_k\| s_k\|,$$

and thus, using (3.9), that

$$\|B_k s_k + \sigma_k \|s_k\| s_k\| \geq (1 - \kappa_\theta) \|g_k\| > (1 - \kappa_\theta) \epsilon.$$

Taking this bound, (2.13), (2.15), (2.2) and **A.4** into account, we deduce that

$$\begin{aligned} (1 - \kappa_\theta) \epsilon &< \kappa_B \|s_k\| + \sigma_k \|s_k\|^2 \\ &\leq \left\{ \kappa_B + 3 \max \left[ \|B_k\|, \sqrt{\sigma_k \|g_k\|} \right] \right\} \|s_k\| \\ &\leq \left\{ \kappa_B + 3 \max \left[ \kappa_B, \sqrt{\sigma_k \kappa_{\text{ubg}}} \right] \right\} \|s_k\|, \end{aligned}$$

proving (3.7).  $\square$

We are now able to deduce that the inner loop of Algorithm ARC-FDH terminates in a bounded number of iterations and hence that the desired accuracy on the Hessian approximation is obtained.

**Lemma 3.2.** *Suppose that **A.1**, **A.4** and (2.21) hold. Then the total number of times where a return from Step 4 to Step 1 is executed in Algorithm ARC-FDH is bounded above by*

$$\left\lceil \frac{\log \kappa_h + \frac{3}{2} \log \epsilon}{\log \gamma_3} \right\rceil_+ \quad (3.10)$$

where  $\kappa_h > 0$  is a constant independent of  $n$  and where  $\lceil \alpha \rceil_+$  denotes the maximum of zero and the first integer larger or equal to  $\alpha$ . Moreover **A.3** holds.

**Proof.** The inequality (3.7) and (2.19) give that

$$(1 - \kappa_\theta)\epsilon \leq \max \left[ 4\kappa_B, \kappa_B + 3\sqrt{\frac{\kappa_\sigma \kappa_{\text{ubg}}}{\epsilon}} \right] \|s_k\| \leq \frac{4\kappa_B}{\epsilon^{1/2}} \|s_k\|, \quad (3.11)$$

where we have used the bound  $\kappa_B \geq \sqrt{\kappa_\sigma \kappa_{\text{ubg}}}$  and the inclusion  $\epsilon \in (0, 1)$  to deduce the last inequality. Now the loop between Steps 1 and 4 of Algorithm ARC-FDH terminates as soon as (3.4) is violated, which must happen if  $j$  is large enough to ensure that

$$h_{k,j} = \gamma_3^j h_{k,0} \leq \gamma_3^j \leq \frac{\kappa_{\text{hs}}(1 - \kappa_\theta)}{4\kappa_B} \epsilon^{3/2} \leq \kappa_{\text{hs}} \|s_{k,j}\|, \quad (3.12)$$

where we have successively used the mechanism of the algorithm, and (3.11). The second inequality in (3.12) and the decreasing nature of the sequence  $\{h_k\}$  then ensures that (3.3) must hold for all  $k$  after at most (3.10) (with  $\kappa_h = \kappa_{\text{hs}}(1 - \kappa_\theta)/4\kappa_B$ ) reductions of the stepsize by  $\gamma_3$ , which proves the first part of the lemma. Finally, (3.3) and (3.2) imply also that (2.14) holds for  $B_k$ . This with (2.13) ensures that **A.3** is satisfied.  $\square$

We may then conclude with our main result for this section.

**Theorem 3.3.** *Suppose that **A.1**, **A.2** and **A.4** hold, that  $\epsilon \in (0, 1)$  is given and that (2.21) holds. Then Algorithm ARC-FDH terminates after at most*

$$N_1^s \stackrel{\text{def}}{=} 1 + \lceil \kappa_S^s \epsilon^{-3/2} \rceil, \quad (3.13)$$

successful iterations and at most

$$N_1 \stackrel{\text{def}}{=} \lceil \kappa_S \epsilon^{-3/2} \rceil \quad (3.14)$$

iterations in total, where  $\kappa_S^s$  and  $\kappa_S$  are given by (2.25) and (2.26), respectively. As a consequence, the ARC-FDH algorithm terminates after at most

$$(n + 1)N_1^s + n \left\lceil \frac{\log \kappa_h + \frac{3}{2} \log \epsilon}{\log \gamma_3} \right\rceil_+ \quad (3.15)$$

gradient evaluations and at most  $N_1$  objective function evaluations.

**Proof.** Lemma 3.2 ensures that **A.3** holds. Theorem 2.5 is thus applicable and the number of successful iterations is therefore bounded by (2.23), while the total number of iterations is bounded by (2.24). The bound (3.15) and the bound of the number of function evaluations then follows from Lemma 3.2 and the observation that, in addition to the computation of  $\nabla_x f(x_k)$  (at successful iterations only) and  $f(x_k)$ , each successful iteration involves an estimation of the Hessian by finite differences, each of which requires  $n$  gradient evaluations, plus possibly at most (3.10) additional Hessian estimations at the same cost.  $\square$

Very broadly speaking, we therefore require that at most

$$O\left(n \left[ \left\lceil \frac{1}{\epsilon^{3/2}} \right\rceil + \lceil \lceil \log \epsilon \rceil \rceil \right] \right) \quad (3.16)$$

gradient and

$$O\left(\left\lceil \frac{1}{\epsilon^{3/2}} \right\rceil\right)$$

function evaluations in the worst-case. This is qualitatively very similar to the bound (2.24) for the original ARC algorithm.

We close this section by observing that better bounds may be obtained if we assume that the Hessian has a known sparsity pattern. The finite-difference scheme may then be adapted (see Powell and Toint, 1979, or Goldfarb and Toint, 1984) to require much less than  $n$  gradient differences to obtain a Hessian approximation, in which case the factor  $n$  in (3.16) may often be replaced by a small constant. Similar gains can be obtained if  $f$  is partially separable (Griewank and Toint, 1982). Finally, parallel evaluations of the gradient in Step 1 may also result in substantial computational savings.

## 4 A derivative-free ARC variant

We are now interested in pursuing the same idea further and considering a derivative-free variant of the ARC algorithm, where both gradients and Hessians are approximated by finite differences. However, this introduces two additional difficulties: the approximation techniques used for the gradient and Hessian should be clarified and balanced, and some results we relied on in the previous section (in particular Lemmas 2.2 and 2.3) have to be revisited because they depend on the true gradient of the objective function, which is no longer available.

Consider the approximation of gradients and Hessians first. From the discussion above, we see that preserving (2.14) is necessary for using results for the original ARC algorithm. It is then natural to seek a higher degree of accuracy for the gradient itself, since this is the quantity that the algorithm drives to zero. We therefore suggest using a central difference scheme for the gradient, in which a quotient of the form

$$\frac{f(x_k + t_k e_i) - f(x_k - t_k e_i)}{2t_k} \quad (4.1)$$

for some stepsize  $t_k > 0$  is used to approximate the  $i$ -th component of the gradient at  $x_k$ . It is well-known (see Nocedal and Wright, 1999, Section 7.1) that such a scheme ensures the bound

$$\|\nabla_x f(x_k) - g_k\| \leq \kappa_{\text{egt}} t_k^2 \quad (4.2)$$

for some constant  $\kappa_{\text{egt}} \in [0, L_H]$ , where  $g_k$  is now the vector *approximating*  $\nabla_x f(x_k)$ , i.e. whose  $i$ -th component is given by (4.1). Similarly, we may approximate the  $(i, j)$ -th entry of the Hessian at  $x_k$  by a quotient of the form

$$\frac{f(x_k + t_k e_i + t_k e_j) - f(x_k + t_k e_i) - f(x_k + t_k e_j) + f(x_k)}{t_k^2}, \quad (4.3)$$

(see Nocedal and Wright, 1999, Section 7.1), yielding the error bound

$$\|\nabla_{xx} f(x_k) - B_k\| \leq \kappa_{\text{Ht}} t_k \quad (4.4)$$

for some constant  $\kappa_{\text{Ht}} \in [0, L_H]$ , where  $B_k$  is the symmetric matrix whose  $(i, j)$ -th entry (for  $i \geq j$ ) is given by the quotient (4.3). Note that (4.4) give the same type of error bound as (3.2) above, and we are again interested in an algorithm which guarantees (2.14) from (4.4), i.e. such that

$$t_k \leq \kappa_{\text{ts}} \|s_k\| \quad (4.5)$$

for all  $k \geq 0$  and some constant  $\kappa_{\text{ts}} > 0$ .

The gradient approximation scheme also raises the question of proper termination of any algorithm using  $g_k$  rather than  $\nabla_x f(x_k)$ . Since this latter quantity is unavailable by assumption, it is impossible to test its norm against the threshold  $\epsilon$ . The next best thing is to test  $\|g_k\|$  for a sufficiently small difference stepsize  $t_k$ . More specifically, if

$$\|g_k\| \leq \frac{1}{2}\epsilon \quad \text{and} \quad t_k \leq \sqrt{\frac{\epsilon}{2\kappa_{\text{egt}}}} \stackrel{\text{def}}{=} t_\epsilon \quad (4.6)$$

then (4.2) and the triangle inequality ensure that  $\|\nabla_x f(x_k)\| \leq \epsilon$ , as requested. In what follows, we assume that we know a suitable value for  $\kappa_{\text{egt}}$  or, equivalently, of  $t_\epsilon$ , and then use (4.6) for detecting an approximate first-order critical point. *The worst-case complexity is therefore to be understood as the maximum number of function evaluations necessary for the test (4.6) to hold.*

Using these ideas, we may now state the ARC-DFO variant of the ARC algorithm on the following page.

As was the convention for Algorithm ARC-FDH above, we denote by  $B_k$ ,  $g_k$  and  $g_k^+$  the quantities  $B_{k,j}$ ,  $g_{k,j}$  and  $g_{k,j}^+$  obtained at the end of the loop between Steps 3 and 7 (we show below that this loop terminates finitely). It is also clear that the stepsizes  $t_k$  are monotonically decreasing. We also see that Step 7 ensures (4.5). We next verify that the Hessian approximations remains bounded and that loop between Steps 3 and 7 always terminates after a finite number of iterations.

**Algorithm 4.1: ARC-DFO**

**Step 0:** An initial starting point  $x_0$  is given, as well as a user-defined accuracy threshold  $\epsilon \in (0, 1)$ . Choose a stepsize  $t_{0,0} \leq t_\epsilon$ . Set  $k = 0$  and  $j = 0$ .

**Step 1:** Estimate  $g_{0,0}$  using (4.1) with stepsize  $t_{0,j}$ .

**Step 2:** If  $\|g_{0,j}\| \leq \frac{1}{2}\epsilon$ , terminate with approximate solution  $x_0$ .

**Step 3:** Estimate  $B_{k,j}$  using (4.3) with stepsize  $t_{k,j}$ .

**Step 4:** Compute a step  $s_{k,j}$  satisfying (2.3)–(2.7).

**Step 5:** Estimate  $g_{k,j}^+$  using (4.1) with  $x_k$  replaced by  $x_k + s_{k,j}$  and the stepsize  $t_{k,j}$ .

**Step 6:** If  $\|g_{k,j}^+\| \leq \frac{1}{2}\epsilon$ , terminate with approximate solution  $x_k + s_{k,j}$ .

**Step 7:** If

$$t_{k,j} > \kappa_{\text{ts}} \min[\|s_{k,j}\|, \|g_{k,j}\|] \quad (4.7)$$

set  $t_{k,j+1} = \gamma_3 t_{k,j}$ , increment  $j$  by one and return to Step 3. Otherwise, set  $s_k = s_{k,j}$  and  $t_k = t_{k,j}$ .

**Step 8:** Compute  $f(x_k + s_k)$  and

$$\rho_k = \frac{f(x_k) - f(x_k + s_k)}{-m_k(s_k)}. \quad (4.8)$$

**Step 9:** Set

$$x_{k+1} = \begin{cases} x_k + s_k & \text{if } \rho_k \geq \eta_1, \\ x_k & \text{otherwise,} \end{cases} \quad \text{and} \quad g_{k+1,0} = \begin{cases} g_{k,j}^+ & \text{if } \rho_k \geq \eta_1, \\ g_{k,j} & \text{otherwise.} \end{cases}$$

**Step 10:** Set

$$\sigma_{k+1} \in \begin{cases} (0, \sigma_k] & \text{if } \rho_k > \eta_2, & \text{[very successful iteration]} \\ [\sigma_k, \gamma_1 \sigma_k] & \text{if } \eta_1 \leq \rho_k \leq \eta_2, & \text{[successful iteration]} \\ [\gamma_1 \sigma_k, \gamma_2 \sigma_k] & \text{otherwise.} & \text{[unsuccessful iteration]} \end{cases} \quad (4.9)$$

**Step 11:** Set  $t_{k+1,0} = t_k$  and  $j = 0$ . Increment  $k$  by one and return to Step 3 if  $\rho_k \geq \eta_1$  or to Step 4 otherwise.

**Lemma 4.1.** *Suppose that **A.1** and **A.4** hold. Then there exists constants  $\kappa_B > 1$  and  $\kappa_{ng} > 0$  such that, if  $B_{k,j}$  is estimated at Step 3, then*

$$\|g_k\| \leq \kappa_{ng} \quad \text{and} \quad \|B_k\| \leq \kappa_B. \quad (4.10)$$

for all  $k \geq 0$ . Moreover, we have that

$$\|s_k\| \geq \frac{(1 - \kappa_\theta) \epsilon}{\max [4\kappa_B, \kappa_B + 3\sqrt{\sigma_k \kappa_{ubg}}]} \quad (4.11)$$

and there exists a  $\kappa(\sigma_k) > 0$  such that, at iteration  $k$  of Algorithm ARC-DFO, the loop between Steps 3 and 7 terminates in at most

$$\left\lceil \frac{\log \kappa(\sigma_k) + \log \epsilon}{\log \gamma_3} \right\rceil_+ \quad (4.12)$$

iterations. Finally, the inequalities

$$\|g_k - \nabla_x f(x_k)\| \leq \kappa_{egt} \kappa_{ts} \|s_k\|^2, \quad (4.13)$$

$$\|g_k^+ - \nabla_x f(x_k + s_k)\| \leq \kappa_{egt} \kappa_{ts} \|s_k\|^2 \quad (4.14)$$

and

$$\|B_k - \nabla_{xx} f(x_k)\| \leq \kappa_{eHt} \kappa_{ts} \|s_k\|. \quad (4.15)$$

hold for each  $k \geq 0$ .

**Proof.** Consider iteration  $k$ . As in Lemma 3.2, we obtain that  $\|B_{k,j}\| \leq \kappa_B$  and therefore that the second inequality in (4.10) holds. The proof of the first is similar in spirit:

$$\|g_k\| \leq \|g_k - \nabla_x f(x_k)\| + \|\nabla_x f(x_k)\| \leq \kappa_{egt} + \kappa_{ubg} \stackrel{\text{def}}{=} \kappa_{ng},$$

where we used (4.2), the inequality  $t_{k,j} \leq t_{0,0} \leq 1$  and **A.4**. Observe now that the mechanism of the algorithm implies that, as long as the algorithm isn't terminated,

$$\|g_k\| \geq \frac{1}{2} \epsilon. \quad (4.16)$$

As in the proof of Lemma 3.1 (using (4.16) instead of (3.9)), we may now derive that (4.11) holds for all  $k$ . Defining

$$\mu(\sigma_k) \stackrel{\text{def}}{=} \frac{1 - \kappa_\theta}{\max [4\kappa_B, \kappa_B + 3\sqrt{\sigma_k \kappa_{ubg}}]}$$

this lower bound may then be used to deduce that the loop between Steps 3 and 7 terminates as soon as (4.7) is violated, which must happen if  $j$  is large enough to ensure that

$$t_{k,j} = \gamma_3^j t_{k,0} \leq \gamma_3^j \leq \kappa_{ts} \min [\mu(\sigma_k), \frac{1}{2}] \epsilon \leq \kappa_{ts} \min [\|s_k\|, \|g_k\|], \quad (4.17)$$



where we used (4.16) to derive the last inequality. This implies that  $j$  never exceeds

$$\left\lceil \frac{\log \{[\kappa_{ts} \min [\mu(\sigma_k), \frac{1}{2}]]\} + \log \epsilon}{\log \gamma_3} \right\rceil_+,$$

which in turn yields (4.12) with  $\kappa(\sigma_k) \stackrel{\text{def}}{=} \kappa_{ts} \min [\mu(\sigma_k), \frac{1}{2}]$ . Since the loop between Steps 3 and 7 always terminates finitely, (4.5) holds for all  $k \geq 0$  and the inequalities (4.13)–(4.15) then follow from (4.2) and (4.4).  $\square$

Unfortunately, several of the basic properties of the ARC algorithm mentioned in Section 2 can no longer be deduced from existing theory. This is the case of (2.19), (2.18) and (2.20), which we thus need to reconsider.

The proof of (2.19) is involved and needs to be restarted from the Cauchy condition (2.5)–(2.6). This condition is known to imply the inequality

$$f(x_k) - m_k(s_k) \geq \kappa_C \|g_k\| \min \left[ \frac{\|g_k\|}{1 + \|B_k\|}, \sqrt{\frac{\|g_k\|}{\sigma_k}} \right] \quad (4.18)$$

for some constant  $\kappa_C \in (0, 1)$  (see Lemma 1.1 in Cartis et al., 2009a). We may then build on this relation in the next two useful lemmas inspired from Cartis et al. (2009a).

**Lemma 4.2.** [See Lemmas 3.2 in Cartis et al., 2009a] *Suppose that **A.1** and **A.4** hold, and that*

$$\sqrt{\sigma_k \|s_k\|} \geq \frac{108\sqrt{2}}{1 - \eta_2} (L_g + \kappa_{egt} \kappa_{ts}^2 (\kappa_{ubg} + \kappa_{egt}) + \kappa_B) \stackrel{\text{def}}{=} \kappa_{HB}. \quad (4.19)$$

*Then iteration  $k$  of Algorithm ARC-DFO is very successful and*

$$\sigma_{k+1} \leq \sigma_k. \quad (4.20)$$

**Proof.** From (4.19), we have that  $g_k \neq 0$ , and thus (4.18) implies that  $f(x_k) > m_k(s_k)$ . It then follows from (4.8) that

$$\rho_k > \eta_2 \Leftrightarrow \nu_k \stackrel{\text{def}}{=} f(x_k + s_k) - f(x_k) - \eta_2 [m_k(s_k) - f(x_k)] < 0.$$

We immediately note that, for  $k \geq 0$ ,

$$\nu_k = f(x_k + s_k) - m_k(x_k) + (1 - \eta_2) [m_k(s_k) - f(x_k)].$$

We then develop the first term in the right-hand side of this expression using a Taylor expansion of  $f(x_k + s_k)$ , giving that, for  $k \geq 0$ ,

$$f(x_k + s_k) - m_k(s_k) = \langle \nabla_x f(\xi_k) - g_k, s_k \rangle - \frac{1}{2} \langle s_k, B_k s_k \rangle - \frac{1}{3} \sigma_k \|s_k\|^3 \quad (4.21)$$

for some  $\xi_k$  in the segment  $(x_k, x_k + s_k)$ . But we observe that

$$\begin{aligned}
\|\nabla_x f(\xi_k) - g_k\| &\leq \|\nabla_x f(\xi_k) - \nabla_x f(x_k)\| + \|\nabla_x f(x_k) - g_k\| \\
&\leq L_g \|s_k\| + \kappa_{\text{egt}} t_k^2 \\
&\leq L_g \|s_k\| + \kappa_{\text{egt}} \kappa_{\text{ts}}^2 \|s_k\| \|g_k\| \\
&\leq [L_g + \kappa_{\text{egt}} \kappa_{\text{ts}}^2 (\|\nabla_x f(x_k)\| + \|\nabla_x f(x_k) - g_k\|)] \|s_k\| \\
&\leq [L_g + \kappa_{\text{egt}} \kappa_{\text{ts}}^2 (\kappa_{\text{ubg}} + \kappa_{\text{egt}})] \|s_k\|,
\end{aligned}$$

where we successively used the triangle inequality, (2.11), (4.2), the negation of (4.7), **A.4** and the inequality  $t_k \leq 1$ . Thus the Cauchy-Schwartz inequality, (4.21) and the second inequality of (4.10) give that, for  $k \geq 0$ ,

$$f(x_k + s_k) - m_k(s_k) \leq [L_g + \kappa_{\text{egt}} \kappa_{\text{ts}}^2 (\kappa_{\text{ubg}} + \kappa_{\text{egt}}) + \kappa_{\text{B}}] \|s_k\|^2. \quad (4.22)$$

The proof of the lemma then follows exactly as in Lemma 3.2 in Cartis et al. (2009a), using (4.18), with (4.22) playing the role of inequality (3.9) and  $L_g + \kappa_{\text{egt}} \kappa_{\text{ts}} (\kappa_{\text{ubg}} + \kappa_{\text{egt}})$  playing the role of  $\kappa_{\text{H}}$ .  $\square$

We may then recover boundedness of the regularization parameters.

**Lemma 4.3.** *Suppose that **A.1** and **A.4** hold. Then there exists a  $\kappa_\sigma > 0$  such that (2.17) holds for all  $k \geq 0$ .*

**Proof.** The proof is identical to that of Lemma 3.3 in Cartis et al. (2009a), giving  $\kappa_\sigma \stackrel{\text{def}}{=} \gamma_2 \kappa_{\text{HB}}^2$ .  $\square$

Again, we replace (2.17) by (2.19) and, since  $\kappa_\sigma$  does not depend on  $\kappa_{\text{B}}$ , possibly increase  $\kappa_{\text{B}}$  to ensure that  $\kappa_{\text{B}} \geq \kappa_\sigma \kappa_{\text{ubg}}$  without loss of generality. Armed with these results, we may return to Lemma 4.1 above and obtain stronger conclusions.

**Lemma 4.4.** *Suppose that **A.1** and **A.4** hold. Then there exists a constant  $\kappa_t > 0$  such that the return from Step 7 to Step 3 of Algorithm ARC-DFO can only be executed at most*

$$\left\lceil \frac{\log \kappa_t + \frac{3}{2} \log \epsilon}{\log \gamma_3} \right\rceil_+ \quad (4.23)$$

*times during the entire run of the algorithm.*

**Proof.** Replacing (2.17) into (4.11), we obtain that, for all  $k \geq 0$

$$\|s_k\| \geq \frac{(1 - \kappa_\theta) \epsilon}{\max \left[ 4\kappa_{\text{B}}, \kappa_{\text{B}} + 3\sqrt{\kappa_\sigma \kappa_{\text{ubg}}/\epsilon} \right]} \geq \frac{(1 - \kappa_\theta) \epsilon^{3/2}}{4\kappa_{\text{B}}} \stackrel{\text{def}}{=} \kappa_{\text{sc}} \epsilon^{3/2},$$

Thus no return from Step 7 to Step 3 of Algorithm ARC-DFO is possible as soon as  $j \geq 0$ , the total number of times this return is executed, is large enough to ensure that

$$t_{k,j} = \gamma_3^j t_{0,0} \leq \gamma_3^j \leq \kappa_{\text{ts}} \min \left[ \kappa_{\text{sc}} \epsilon^{3/2}, \frac{1}{2} \epsilon \right] \leq \kappa_{\text{ts}} \min \left[ \|s_{k,j}\|, \|g_{k,j}\| \right],$$

where we have derived the last inequality using the fact that  $\|g_{k,j}\| \geq \frac{1}{2}\epsilon$  as long as the algorithm has not terminated. This imposes that

$$j \leq \frac{1}{\log \gamma_3} \min [\log (\kappa_{ts} \kappa_{s\epsilon}) + \frac{3}{2} \log \epsilon, \log (\frac{1}{2} \kappa_{ts}) + \log \epsilon],$$

and the desired bound on  $j$  follows with  $\kappa_t = \kappa_{ts} \min[\kappa_{s\epsilon}, \frac{1}{2}]$ .  $\square$

We may also revisit the second part of Lemma 2.2 in the derivative-free context. Our proof is directly inspired by Lemma 5.2 in Cartis et al. (2009a).

**Lemma 4.5.** *Suppose that A.1 and A.4 hold. Then there exists a  $\sigma_{\max} > 0$  independent of  $\epsilon$  such that (2.18) holds for all  $k \geq 0$ .*

**Proof.** Using (2.1), the Cauchy-Schwarz and the triangle inequalities, (4.13), (2.12) and (4.15), we know that

$$\begin{aligned} f(x_k + s_k) - m_k(s_k) &\leq \|\nabla_x f(x_k) - g_k\| \|s_k\| \\ &\quad + \frac{1}{2} [\|\nabla_{xx} f(\xi_k) - \nabla_{xx} f(x_k)\| + \|\nabla_{xx} f(x_k) - B_k\|] \|s_k\|^2 \\ &\quad - \frac{1}{3} \sigma_k \|s_k\|^3 \\ &\leq [\kappa_{\text{egt}} \kappa_{ts} + \frac{1}{2}(L_H + \kappa_{\text{eHt}} \kappa_{ts}) - \frac{1}{3} \sigma_k] \|s_k\|^3 \end{aligned}$$

for some  $\xi_k \in [x_k, x_k + s_k]$ . Thus, using (4.8) and (2.16),

$$\rho_k - 1 = \frac{f(x_k + s_k) - m_k(s_k)}{-m_k(s_k)} \leq \frac{\kappa_{\text{egt}} \kappa_{ts} + \frac{1}{2}(L_H + \kappa_{\text{eHt}} \kappa_{ts}) - \frac{1}{3} \sigma_k}{\frac{1}{6} \sigma_k} \leq 1 - \eta_2$$

as soon as

$$\sigma_k \geq \frac{2\kappa_{\text{egt}} \kappa_{ts} + L_H + \kappa_{\text{eHt}} \kappa_{ts}}{1 - \frac{1}{3} \eta_2}.$$

As a consequence, iteration  $k$  is then very successful and  $\sigma_{k+1} \leq \sigma_k$ . It then follows that (2.18) holds with

$$\sigma_{\max} = \max \left[ \sigma_0, \frac{\gamma_2(2\kappa_{\text{egt}} \kappa_{ts} + L_H + \kappa_{\text{eHt}} \kappa_{ts})}{1 - \frac{1}{3} \eta_2} \right].$$

$\square$

It then remains to show that, under (4.13)–(4.15), an analog of Lemma 2.3 holds for the derivative-free case.

**Lemma 4.6.** *Suppose that A.1 and A.4 hold. Then there exists a constant  $\kappa_g > 0$  such that, for all  $k \geq 0$ ,*

$$\|s_k\| \geq \kappa_g \sqrt{\|g_k^+\|}. \quad (4.24)$$

**Proof.** We first observe, using the triangle inequality, (4.14) and (2.7), that

$$\begin{aligned} \|g_k^+\| &\leq \|g_k^+ - \nabla_x f(x_k + s_k)\| + \|\nabla_x f(x_k + s_k) - \nabla_x m_k(s_k)\| + \|\nabla_x m_k(s_k)\| \\ &\leq \kappa_{\text{egt}}\kappa_{\text{ts}}\|s_k\|^2 + \|\nabla_x f(x_k + s_k) - \nabla_x m_k(s_k)\| + \kappa_\theta \min[1, \|s_k\|] \|g_k\|. \end{aligned} \quad (4.25)$$

for all  $k \geq 0$ . The second term on this last right-hand side may then be bounded for all  $k \geq 0$  by

$$\begin{aligned} \|\nabla_x f(x_k + s_k) - \nabla_x m_k(s_k)\| &\leq \|\nabla_x f(x_k) - g_k\| + \left\| \int_0^1 [\nabla_{xx}(x_k + \alpha s_k) - B_k] s_k d\alpha \right\| + \sigma_k \|s_k\|^2 \\ &\leq \left\| \int_0^1 \{[\nabla_{xx}(x_k + \alpha s_k) - \nabla_{xx}f(x_k)] + [\nabla_{xx}f(x_k) - B_k]\} s_k d\alpha \right\| \\ &\quad + \|\nabla_x f(x_k) - g_k\| + \sigma_k \|s_k\|^2 \\ &\leq \max_{\alpha \in [0,1]} \|\nabla_{xx}(x_k + \alpha s_k) - \nabla_{xx}f(x_k)\| \|s_k\| \\ &\quad + (\kappa_{\text{eHt}} + \kappa_{\text{egt}})\kappa_{\text{ts}}\|s_k\|^2 + \sigma_{\text{max}}\|s_k\|^2 \\ &\leq [L_H + (\kappa_{\text{eHt}} + \kappa_{\text{egt}})\kappa_{\text{ts}} + \sigma_{\text{max}}]\|s_k\|^2, \end{aligned} \quad (4.26)$$

where we successively used the mean-value theorem, (2.1), the triangle inequality, (2.12), (4.13), (4.15) and (2.18). We also have, using the triangle inequality, (4.13), (2.11) and (4.14), that

$$\begin{aligned} \|g_k\| &\leq \|g_k - \nabla_x f(x_k)\| + \|\nabla_x f(x_k)\| \\ &\leq \kappa_{\text{egt}}\kappa_{\text{ts}}\|s_k\|^2 + \|\nabla_x f(x_k + s_k)\| + L_g\|s_k\| \\ &\leq \kappa_{\text{egt}}\kappa_{\text{ts}}\|s_k\|^2 + \|\nabla_x f(x_k + s_k) - g_k^+\| + \|g_k^+\| + L_g\|s_k\| \\ &\leq 2\kappa_{\text{egt}}\kappa_{\text{ts}}\|s_k\|^2 + \|g_k^+\| + L_g\|s_k\|. \end{aligned}$$

which implies that, for all  $k \geq 0$ ,

$$\kappa_\theta \min[1, \|s_k\|] \|g_k\| \leq (2\kappa_\theta\kappa_{\text{egt}}\kappa_{\text{ts}} + \kappa_\theta L_g)\|s_k\|^2 + \kappa_\theta \|g_k^+\|. \quad (4.27)$$

Therefore, substituting (4.26) and (4.27) into (4.25), we obtain that, for all  $k \geq 0$ ,

$$\|g_k^+\| \leq \kappa_{\text{egt}}\kappa_{\text{ts}}\|s_k\|^2 + [L_H + (\kappa_{\text{eHt}} + \kappa_{\text{egt}})\kappa_{\text{ts}} + \sigma_{\text{max}}]\|s_k\|^2 + (2\kappa_\theta\kappa_{\text{egt}}\kappa_{\text{ts}} + \kappa_\theta L_g)\|s_k\|^2 + \kappa_\theta \|g_k^+\|.$$

and thus

$$(1 - \kappa_\theta)\|g_k^+\| \leq [\kappa_\theta L_g + L_H + \kappa_{\text{ts}}(\kappa_{\text{eHt}} + 2\kappa_{\text{egt}}(1 + \kappa_\theta)) + \sigma_{\text{max}}]\|s_k\|^2$$

for all  $k \geq 0$ . This gives (4.24) with

$$\kappa_g \stackrel{\text{def}}{=} \sqrt{\frac{1 - \kappa_\theta}{\kappa_\theta L_g(1 + \kappa_\theta) + L_H + \kappa_{\text{ts}}(\kappa_{\text{eHt}} + 2\kappa_{\text{egt}}(1 + \kappa_\theta)) + \sigma_{\text{max}}}}.$$

□

We are thus in principle again in position to apply the oracle complexity results for the ARC algorithm. Unfortunately, Theorem 2.5 may no longer be applied as such (as it requires the true gradient of the objective function), but our final theorem is derived in a very similar manner.

**Theorem 4.7.** *Suppose that **A.1**, **A.2** and **A.4** hold, that  $\epsilon \in (0, 1)$  is given and that (2.21) holds. Then Algorithm ARC-DFO terminates after at most*

$$N_1^s \stackrel{\text{def}}{=} 1 + \lceil \kappa_S^s \epsilon^{-3/2} \rceil, \quad (4.28)$$

*successful iterations and at most*

$$N_1 \stackrel{\text{def}}{=} \lceil \kappa_S \epsilon^{-3/2} \rceil \quad (4.29)$$

*iterations in total, where  $\kappa_S^s$  and  $\kappa_S$  are given by (2.25) and (2.26), respectively. As a consequence, the ARC algorithm terminates after at most*

$$(N_1 - N_1^s)(1 + 2n) + N_1^s \left[ \frac{n^2 + 5n + 2}{2} \right] + \left[ \frac{n^2 + 3n}{2} \right] \left\lceil \frac{\log \kappa_t + \frac{3}{2} \log \epsilon}{\log \gamma_3} \right\rceil_+. \quad (4.30)$$

*objective function evaluations.*

**Proof.** Let

$$\mathcal{K}_\epsilon = \{k \geq 0 \mid \min[\|g_k\|, \|g_{k+1}\|] \geq \frac{1}{2}\epsilon\}.$$

We then deduce from the definition of successful iterations, (2.16) and (4.24) that

$$f(x_k) - f(x_{k+1}) \geq -\eta_1 m_k(s_k) \geq \frac{1}{48} \sigma_{\min} \eta_1 \kappa_g^3 \epsilon^{3/2} \quad \text{for all } k \in \mathcal{K}_\epsilon \cap \mathcal{S}_{k+1}.$$

The mechanism of Algorithm ARC-DFO ensures that the iterates remains unchanged at unsuccessful iterations. If Algorithm ARC-DFO does not terminate before or at iteration  $k$ , we have that  $\mathcal{K}_\epsilon \cap \mathcal{S}_{k+1} = \mathcal{S}_{k+1}$ . Summing up to iteration  $k$ , we therefore obtain that

$$f(x_0) - f(x_{k+1}) \geq \sum_{i \in \mathcal{S}_j} [f(x_i) - f(x_{i+1})] \geq \frac{1}{48} \sigma_{\min} \eta_1 \kappa_g^3 \epsilon^{3/2} |\mathcal{S}_{k+1}|$$

Using now **AS.2**, we conclude that

$$|\mathcal{S}_{k+1}| \leq \frac{48(f(x_0) - f_{\text{low}})}{\sigma_{\min} \eta_1 \kappa_g^3 \epsilon^{3/2}},$$

from which (4.28) follows with

$$\kappa_S^s = \frac{48(f(x_0) - f_{\text{low}})}{\sigma_{\min} \eta_1 \kappa_g^3}.$$

We then use Lemma 2.4 to deduce (4.29). In we ignore the estimations of  $B_{k,j}$  in Step 3 after a return from Step 7, we now observe that each successful iteration involves up to

$$1 + 2n + \left( \frac{n(n+1)}{2} \right)$$

function evaluations, while unsuccessful iterations involves  $1 + 2n$  evaluations. Adding the two, we obtain a number of

$$(N_1 - N_1^s)(1 + 2n) + N_1^s \left[ 1 + 2n + \frac{n(n+1)}{2} \right]$$

evaluations at most, to which we have to add those needed in the loop between Steps 3 and 7, whose number does not exceed

$$\left[ n + \frac{n(n+1)}{2} \right] \left[ \frac{\log \kappa_t + \frac{3}{2} \log \epsilon}{\log \gamma_3} \right]_+.$$

The resulting grand total is then given by (4.30).  $\square$

We may again considerably simplify this result (at the cost of a weaker bound). If we assume that the terms in  $n^2$  and  $n$  dominate the constants, we obtain that, in the worst case, at most

$$O \left( \frac{n^2 + 5n}{2} \left[ 1 + \lceil \log \epsilon \rceil_+ + \left[ \frac{1}{\epsilon^{3/2}} \right]_+ \right] \right) \quad (4.31)$$

function evaluations are needed by the ARC-DFO algorithm to achieve approximate criticality in the sense of (4.6). Again, known sparsity of the Hessian or partial separability may reduce the factor  $n^2$  in (4.31) to (typically) a small multiple of  $n$ , thereby bridging the gap between ARC-DFO and ARC itself. The potential benefits of using parallel evaluations of the objective function are even more obvious here than for Algorithm ARC-FDH. Finally notice that automatic differentiation may often be an alternative to derivative-free technology when the source code for the evaluation of  $f$  is available, in which case Algorithm ARC-FDH is the natural choice.

## 5 Discussion and conclusions

Comparing algorithms on the basis of their worst-case complexity is always an exercise whose interest is mostly theoretical, but this is especially the case for what we have presented above. Indeed, several factors limit the predictive nature of these results on the practical behaviour of the considered minimization methods. The first is obviously the worst-case nature of the efficiency estimates, which (fortunately) can be quite pessimistic in view of expected or observed efficiency. The second, which is specific to the results presented here, is the intrinsic limitations induced by the use of finite-precision arithmetic. In the context of actual computation, not only it is unrealistic to consider vanishingly

small values of  $\epsilon$ , but the choice of arbitrarily small finite-differences stepsizes is also very questionable<sup>4</sup>, even if difficulties caused by finite precision may be attenuated by using multiple-precision packages. The following comments should therefore be considered as interesting theoretical considerations throwing some light on the fundamental differences between algorithms, even if their practical relevance to actual numerical performance is potentially remote. Designing and studying worst-case analysis in the presence of round-off errors remains an interesting challenge.

We first note that the gap in worst-case performance between second-order (ARC), first-order (ARC-FDH) and derivative-free (ARC-DFO) methods is remarkably small if one consider the associated bounds in the asymptotic regime where  $\epsilon$  tends to zero. The effect of finite-difference schemes is, up to constants, limited to the occurrence of a multiplicative factor of size  $1 + |\log \epsilon|$ , which may be considered as modest. The more significant effect is not depending on the  $\epsilon$ -asymptotics, but rather depending on the dimension  $n$  of the problem: as expected, derivative-free methods suffer most in this respect, with bounds depending on  $n^2$  rather than  $n$  for first-order methods or a constant for second-order ones.

The bounds for derivative-free methods are also interesting to compare with those derived by Vicente (2010), where direct-search type methods are shown to require at most  $O(\epsilon^{-2})$  iterations to find a point  $x_k$  satisfying  $\|\nabla_x f(x_k)\| \leq \epsilon$  when applied to function with Lipschitz continuous gradients<sup>5</sup>. At iteration  $k$ , such methods compute the function values  $\{f(x_k + \alpha_k d \mid d \in \mathcal{D}_k)\}$ , where  $\mathcal{D}_k$  is a positive spanning set for  $\mathbb{R}^n$  and  $\alpha_k$  an iteration-dependent stepsize. If one of these value is (sufficiently) lower than  $f(x_k)$  the corresponding  $x_k + \alpha_k d$  is chosen as the next iterate and a new iteration started. In the worst-case, an algorithm of this type therefore requires  $n + 1$ <sup>6</sup> function evaluations, and thus its function-evaluation complexity is

$$O\left(n \left\lceil \frac{1}{\epsilon^2} \right\rceil\right)$$

Thus the ARC-DFO algorithm is more advantageous than such direct-search methods (in the worst-case and up to a constant factor) if

$$(n^2 + 5n) \left\lceil \frac{1 + |\log \epsilon|}{\epsilon^{3/2}} \right\rceil = O\left(n \left\lceil \frac{1}{\epsilon^2} \right\rceil\right)$$

that is if

$$n = O\left(\frac{1}{[1 + |\log \epsilon|]\sqrt{\epsilon}}\right).$$

---

<sup>4</sup>Recommended values for these stepsizes are bounded below by adequate roots of machine precision (see Section 8.4.3 in Conn, Gould and Toint, 2000 or Sections 5.4 and 5.6 in Dennis and Schnabel, 1983, for instance).

<sup>5</sup>Note that this inequality cannot be used as a stopping criterion, because  $\nabla_x f(x_k)$  is unknown. The complexity result in Vicente (2010) therefore does not directly indicate how many iterations will be performed by the algorithm before its stopping criterion is activated.

<sup>6</sup>The minimal size of a positive spanning set in  $\mathbb{R}^n$ .

It is interesting to note that this inequality only holds for relatively small  $n$ , especially for values of  $\epsilon$  that are only moderately small, and for a more restrictive class of functions (A.1 vs. Lipschitz gradients). Direct-search methods are thus very often more efficient (in this theoretical sense) than Algorithm ARC-DFO, even if the latter dominates for small  $\epsilon$ .

Finally notice that the central properties needed for proving the complexity result for the ARC-DFO algorithm are the bounds (4.13)–(4.15). These could as well be guaranteed by more sophisticated derivative-free techniques where multivariate interpolation is used to construct Hessian approximation from past points in a suitable neighbourhood of the current iterate (see Conn, Scheinberg and Vicente, 2008, Fasano, Nocedal and Morales, 2009, or Scheinberg and Toint, 2010, for instance). This suggests that a worst-case analysis of these methods might be quite close to that of Algorithm ARC-DFO. Indeed, if gains in the number of function evaluations might be possible by the re-use of these past points compared to using fresh evaluations for establishing a local quadratic model at every iteration, it is not clear that these gains can always be achieved, in particular if every step is large compared the necessary finite-difference stepsize.

## Acknowledgements

The authors are grateful to the Royal Society for its support through the International Joint Project 14265.

## References

- A. Agarwal, P. L. Bartlett, P. Ravikummar, and M. J. Wainwright. Information-theoretic lower bounds on the oracle complexity of convex optimization. *in* ‘Proceedings of the 23rd Annual Conference on Neural Information Processing Systems’, 2009.
- C. Cartis, N. I. M. Gould, and Ph. L. Toint. Adaptive cubic overestimation methods for unconstrained optimization. Part I: motivation, convergence and numerical results. *Mathematical Programming, Series A*, 2009a. DOI: 10.1007/s10107-009-0286-5, 51 pages.
- C. Cartis, N. I. M. Gould, and Ph. L. Toint. An adaptive cubic regularization algorithm for nonconvex optimization with convex constraints and its function-evaluation complexity. Technical Report 08/05R, Department of Mathematics, FUNDP - University of Namur, Namur, Belgium, 2009b.
- C. Cartis, N. I. M. Gould, and Ph. L. Toint. Trust-region and other regularisation of linear least-squares problems. *BIT*, **49**(1), 21–53, 2009c.
- C. Cartis, N. I. M. Gould, and Ph. L. Toint. Adaptive cubic overestimation methods for unconstrained optimization. Part II: worst-case function-evaluation complexity. *Mathematical Programming, Series A*, 2010a. DOI: 10.1007/s10107-009-0337-y.



- C. Cartis, N. I. M. Gould, and Ph. L. Toint. On the complexity of steepest descent, Newton's and regularized Newton's methods for nonconvex unconstrained optimization. *SIAM Journal on Optimization*, **20**(6), 2833–2852, 2010*b*.
- C. Cartis, N. I. M. Gould, and Ph. L. Toint. Worst-case complexity of second-order methods for unconstrained nonconvex optimization. Technical Report (in preparation), Department of Mathematics, FUNDP - University of Namur, Namur, Belgium, 2010*c*.
- A. R. Conn, N. I. M. Gould, and Ph. L. Toint. *Trust-Region Methods*. Number 01 in 'MPS-SIAM Series on Optimization'. SIAM, Philadelphia, USA, 2000.
- A. R. Conn, K. Scheinberg, and L. N. Vicente. *Introduction to Derivative-free Optimization*. MPS-SIAM Optimization series. SIAM, Philadelphia, USA, 2008.
- J. E. Dennis and R. B. Schnabel. *Numerical Methods for Unconstrained Optimization and Nonlinear Equations*. Prentice-Hall, Englewood Cliffs, NJ, USA, 1983. Reprinted as *Classics in Applied Mathematics 16*, SIAM, Philadelphia, USA, 1996.
- G. Fasano, J. Nocedal, and J.-L. Morales. On the geometry phase in model-based algorithms for derivative-free optimization. *Optimization Methods and Software*, to appear, 2009.
- D. Goldfarb and Ph. L. Toint. Optimal estimation of Jacobian and Hessian matrices that arise in finite difference calculations. *Mathematics of Computation*, **43**(167), 69–88, 1984.
- S. Gratton, A. Sartenaer, and Ph. L. Toint. Recursive trust-region methods for multiscale nonlinear optimization. *SIAM Journal on Optimization*, **19**(1), 414–444, 2008.
- A. Griewank. The modification of Newton's method for unconstrained optimization by bounding cubic terms. Technical Report NA/12, Department of Applied Mathematics and Theoretical Physics, University of Cambridge, Cambridge, United Kingdom, 1981.
- A. Griewank and Ph. L. Toint. On the unconstrained optimization of partially separable functions. in M. J. D. Powell, ed., 'Nonlinear Optimization 1981', pp. 301–312, London, 1982. Academic Press.
- A. S. Nemirovski. Efficient methods in convex programming. Lectures notes (online) available on [http://www2.isye.gatech.edu/~nemirovs/OPTI\\_LectureNotes.pdf](http://www2.isye.gatech.edu/~nemirovs/OPTI_LectureNotes.pdf), 1994.
- A. S. Nemirovski and D. B. Yudin. *Problem Complexity and Method Efficiency in Optimization*. J. Wiley and Sons, Chichester, England, 1983.
- Yu. Nesterov. *Introductory Lectures on Convex Optimization*. Applied Optimization. Kluwer Academic Publishers, Dordrecht, The Netherlands, 2004.
- Yu. Nesterov. Accelerating the cubic regularization of Newton's method on convex problems. *Mathematical Programming, Series A*, **112**(1), 159–181, 2008.

- Yu. Nesterov and B. T. Polyak. Cubic regularization of Newton method and its global performance. *Mathematical Programming*, **108**(1), 177–205, 2006.
- J. Nocedal and S. J. Wright. *Numerical Optimization*. Series in Operations Research. Springer Verlag, Heidelberg, Berlin, New York, 1999.
- M. J. D. Powell and Ph. L. Toint. On the estimation of sparse Hessian matrices. *SIAM Journal on Numerical Analysis*, **16**(6), 1060–1074, 1979.
- K. Scheinberg and Ph. L. Toint. Self-correcting geometry in model-based algorithms for derivative-free unconstrained optimization. *SIAM Journal on Optimization*, to appear, 2010.
- S. A. Vavasis. Approximation algorithms for indefinite quadratic programming. *Mathematical Programming*, **57**(2), 279–311, 1992a.
- S. A. Vavasis. *Nonlinear Optimization: Complexity Issues*. International Series of Monographs on Computer Science. Oxford University Press, Oxford, England, 1992b.
- S. A. Vavasis. Black-box complexity of local minimization. *SIAM Journal on Optimization*, **3**(1), 60–80, 1993.
- L. N. Vicente. Worst case complexity of direct search. Technical report, Department of Mathematics, University of Coimbra, Coimbra, Portugal, May 2010.
- M. Weiser, P. Deuffhard, and B. Erdmann. Affine conjugate adaptive Newton methods for nonlinear elastomechanics. *Optimization Methods and Software*, **22**(3), 413–431, 2007.